

## II. REMARKS

### **Formal Matters**

Claims 42-82 are pending after entry of the amendments set forth herein.

Claims 42-82 were examined. Claims 42-49 and 53-57 were rejected. Claims 50-52 and 58-82 were withdrawn from consideration.

Claims 42, 46 and 49 are amended. The amendments to the claims were made solely in the interest of expediting prosecution, and are not to be construed as an acquiescence to any objection or rejection of any claim. Support for the amendments to claims 42, 46, and 49 is found in the claims as originally filed, and throughout the specification, in particular at the following exemplary locations: Specification pages 8, 15, 19, and newly renumbered Figure 6B lanes 2 and 3. Accordingly, no new matter is added by these amendments.

New claims 83 – 100 have been added. Support for new claims 83 – 100 is found in the claims as originally filed, and throughout the specification, in particular at the following exemplary locations: Claims 83 – 91: Specification pages 8, 16, 18, 19, 20, and newly renumbered Figure 6B lane 1; Claims 92-100: Specification pages 8, 15, 17, and 21. Accordingly, no new matter is added by these new claims.

The disclosure has been amended in the specification to address objections noted in the Office Action. The specification has been amended on pages 7 and 10. Support for the amended material can be found in the claims as originally filed, and throughout the specification, in particular at the following exemplary locations: page 7: original claim 1; page 10: original claim 17.

The drawings have been amended to address objections noted in the Office Action. Original Figure 5B and original Figure 6B have been deleted. Original Figures 3C, 3D, 5C and 6C have been amended with revised figure numbers with replacement drawing sheets submitted herewith. The specification has also been amended to remove reference to the deleted figures.

Applicants respectfully request reconsideration of the application in view of the remarks made herein.

### **Examiner Interview**

The undersigned Applicants' representative thanks Examiner Duffy for the courtesy of an in-person interview which took place on November 24, 2003, and which was attended by Examiner Duffy and Applicants' representatives Paula A. Borden and Edward J. Baba.

During the interview, the rejection of claims 42-49 and 53-57 under 35 U.S.C. §112, §102, and §103, was discussed. The amendments to the claims reflect the discussions, which took place during the interview.

### **Drawings**

The drawings filed on 08/02/1999 have been objected to. Replacement figures have been submitted herewith, which replacement figures comply with the requirements for formal drawings. Withdrawal of these objections is respectfully requested.

Original Figures 3C and 3D have been amended to comply with the rules. Original Figure 3C has been amended to revise the figure numbers to Figures 3C-3Y. Accordingly, original Figure 3D has been amended to Figure 3Y, to reflect the renumbering of original Figure 3C.

Original Figures 5B and 6B have been deleted. Original Figures 5C and 6C have been amended to revise the figure numbers to reflect the deletion of Figures 5B and 6B. Replacement drawing sheets are provided herewith. The specification has also been amended to remove reference to the deleted figures.

### **Specification Objections**

The disclosure was objected to because the text references claim numbers. The specification has been amended to remove the reference to claim numbers and alternatively, incorporate the language from the referenced claims into the text of the disclosure. The inserted material corresponds exactly to the text of the original claims that were referenced. Accordingly, no new matter has been added. Therefore, the Examiner is respectfully requested to withdraw the objection.

**Claim objections**

Claims 47, 48, and 49 were objected to under 37 C.F.R. 1.75(c) as allegedly being in improper dependent form for failing to further limit the subject matter of a previous claim.

Without conceding as to the correctness of this rejection, claims 47 and 48 have been canceled rendering the objection of these claims moot. In addition, claim 49 has been amended to remove reference to specific fragments of gp190/MSP1. Therefore, the Examiner is respectfully requested to withdraw this objection.

**Rejection under 35 U.S.C. §112, first paragraph**

***New Matter***

Claims 42-49 and 53-57 were rejected under 35 U.S.C. §112, first paragraph, as allegedly containing subject matter which was not described in the specification in such a way as to reasonably convey to one skilled in the relevant art that the inventors, at the time the application was filed, had possession of the claimed invention.

Specifically, the Office Action stated that the claims read on a reduction of any AT content in any MSP1 nucleotide sequence as compared to any other naturally occurring sequence. As suggested by the Examiner, independent claim 42 has been amended to recite “corresponding naturally occurring” in order to provide a reference point for comparison of a reduced AT content.

The Office Action also states that claims 47 and 48 stand rejected because the claims are drawn to a method of producing a complete gp190/MSP1 polypeptide wherein the nucleotide sequence further comprises an attachment signal or further comprises a signal peptide. Without conceding as to the correctness of this rejection, claims 47 and 48 have been canceled and new claims 83 to 100 have been added in their place. New claims 83 to 100 are directed to two other variations of the gp190/MSP1 protein where either the gp190/MSP1 lacks an attachment signal (claims 83-91) or it lack both an attachment signal and signal peptide (claims 92-100). Support for new claims 83 – 100 can be found in the claims as originally filed, and throughout the specification, in particular at the following exemplary

locations: Claims 83 – 91: Specification pages 8, 16, 18, 19, 20, and Figure 6C lane 1 (renumbered as Figure 6B); Claims 92-100: Specification pages 8, 15, 17, and 21. A schematic representation of examples of polypeptides recited in the claims is provided in the attached Exhibit 1, which was also made available to the examiner prior to the telephone interview on November 24, 2003. Accordingly, no new matter is added by these new claims.

Applicants submit that the new matter rejection of claims 42-49 and 53-57 under 35 U.S.C. §112, first paragraph, has been adequately addressed in view of the remarks set forth above. The Examiner is thus respectfully requested to withdraw the rejection.

### ***Written Description***

Claims 42-49 and 53-57 were rejected under 35 U.S.C. §112, first paragraph, as allegedly lacking written description in the specification. Specifically, the Examiner states that the specification “fails to describe the complete nucleotide sequences encoding naturally occurring gp190/MSP1 proteins corresponding to a representative number of these species, sufficient to describe the genus of nucleotide sequences that are modified to produce nucleotide sequences that are ‘reduced’ in their adenine-thymine content.” In view of the remarks made below, applicants respectfully traverse this rejection.

Under MPEP § 2163.02, the standard for determining compliance with the Written Description requirement is whether the “specification conveys with reasonable clarity to those skilled in the art that, as of the filing date sought, applicant was in possession of the invention as now claimed.” See, e.g., Vas-Cath, Inc. v. Mahurkar, 935 F.2d 1555, 1563-64, 19 USPQ2d 1111, 1117 (Fed. Cir. 1991). Essentially, the specification must “clearly allow persons of ordinary skill in the art to recognize that they invented what is claimed.” See Vas-Cath, 935 F.2d at 1116.

In rejecting the claims the Office Action states that a representative number of nucleic acid sequences have not been provided for other *Plasmodium* species, therefore a skilled artisan could not “envision the detailed chemical structure of the encompassed nucleotide sequences that are used to produce the undescribed proteins of other at least 100 *Plasmodium* species and therefore conception is not achieved until reduction to practice has occurred...the nucleic acid itself is required.” During the November 24, 2003 telephone interview, the Examiner indicated that if the nucleic acid sequence for

MSP-1 of a representative number of species of *Plasmodium* were known at the time the instant application was filed, such would be sufficient to overcome the written description rejection.

The Applicants submit that the nucleic acid sequences for the MSP-1 protein of a representative number of species of *Plasmodium* were known at the time the present application was filed. For example, Chang et al., *Exp. Parasitol.* 67(1):1-11 (1988) (Exhibit 2) discloses the nucleic acid encoding MSP-1 of *Plasmodium Falciparum* (Uganda-Palo Alto strain); Lewis et al., *Mol. Biochem. Parasitol.* 36(3):271-282 (1989) (Exhibit 3) discloses the nucleic acid encoding MSP-1 of *Plasmodium Yoelii*; Deleersnijder et al., *Mol. Biochem. Parasitol.* 43(2):231-244 (1990) (Exhibit 4) discloses the nucleic acid encoding MSP-1 of *Plasmodium Chaubaudi*; Del Portillo et al., *Proc. Natl. Acad. Sci.* 88:4030-4034 (1991) (Exhibit 5) discloses the nucleic acid encoding MSP-1 of *Plasmodium Vivax* (Belum strain); and Gibson et al., *Mol. Biochem. Parasitol.* 50(2):325-333 (1992) (Exhibit 6) closes the nucleic acid encoding MSP-1 of *Plasmodium Vivax* (Sal-1 strain). The Applicants note that the term Merzoite Surface Protein 1 (MSP1 or MSP-1) is also referred to in the literature as: Merzoite Surface Antigen 1 (MSA1 or MSA-1); *Plasmodium* Major Merzoite Surface Antigen (PMMSA); and Major Merzoite Surface Protein Precursor.

Since the sequence of the of MSP-1 gene of various species of *Plasmodium* were available, the methods disclosed in the present application could be readily applied by one skilled in the relevant art of molecular biology to these other MSP-1 genes to produce nucleotide sequences that are “reduced” in their adenine-thymine content. Accordingly, the Applicants submit that the written description rejection of claims 42-49 and 53-57 under 35 U.S.C. § 112, first paragraph, has been adequately addressed in view of the remarks set forth above. The Examiner is thus respectfully requested to withdraw the rejection.

### ***Enablement***

Claim 53 was rejected under 35 U.S.C. § 112, first paragraph, as allegedly containing subject matter which was not described in the specification in such a way as to enable one skilled in the art to which it pertains, or with which it is most nearly connected, to make and/or use in the invention. Without conceding to the correctness of this rejection, claim 53 has been canceled in the spirit of expediting prosecution. Accordingly, this rejection is rendered moot and the Examiner is thus respectfully requested to withdraw the rejection

**Rejection under 35 U.S.C. §112, second paragraph**

Claims 42-49 and 53-57 were rejected under 35 U.S.C. §112, second paragraph, as allegedly indefinite.

The Office Action stated that claim 42 presents a comparison between two sequences that are not structurally defined and therefore, the metes and bounds of the claim cannot be ascertained. Specifically, the Office Action notes that the term “complete” and the terms “naturally occurring nucleotide sequence” do not help to structurally define the two sequences. In the spirit of expediting prosecution, and without conceding as to the correctness of this rejection, claim 42 has been amended to remove the term “complete” and in its place add “having an approximate weight of 190 kD” in order to describe the gp190/MSP1 protein. In addition the term “corresponding” has been added to define the naturally occurring sequence. These amendments were discussed during the November 24, 2003 telephone interview. Support for the amendments of claim 42 can be found in the claims as originally filed, and throughout the specification, in particular at the following exemplary locations: pages 8, 15, 19, and Figure 6C lanes 2 and 3 (renumbered as Figure 6B). The amendments of claim 42 have also been incorporated in newly presented independent claims 83 and 92.

Applicants submit that the rejection of claims 42-49 and 53-57 under 35 U.S.C. §112, second paragraph, has been adequately addressed in view of the remarks set forth above. The Examiner is thus respectfully requested to withdraw the rejection.

**Rejection under 35 U.S.C. §102/103**

Claims 42-49 and 53-57 were rejected under 35 U.S.C. §102(b) or §103 as allegedly unpatentable over Holder et al. ((1985) *Nature* 317:270-273; hereinafter “Holder”).

The Office Action maintained the rejection from the Office Action dated November 20, 2001, and stated that Holder et al. teaches the production of specific fragments of the full length gp190/MSP1 from *P. falciparum*. The Examiner notes that the definition of the term “complete” on page 6 of the specification is inclusive of shorter forms, and that the claim does not define the specific sequence for comparison of the naturally occurring sequence.

As noted above, Claim 42 has been amended to remove the term “complete” and in its place add “having an approximate weight of 190 kD” in order to describe the gp190/MSP1 protein and the term “corresponding” has been added to define the naturally occurring sequence.

Holder does not render claims 42-49 and 53-57 obvious, as there is no mention in Holder of a method for producing gp190/MSP1 having an approximate molecular weight of 190 kD, much less a method of producing gp190/MSP1, comprising expressing a nucleotide sequence encoding gp190/MSP1 in a single expression vector. As stated in the specification, until the instant invention, there was not any successful cloning of the coding region for gp190/MSP1 having an approximate molecular weight of 190 kD. Holder does not disclose a method for solving this problem, nor does Holder suggest any such method. Accordingly, Holder cannot render the instant method as claimed obvious.

Accordingly, the Applicants submit that the rejection of claims 42-49 and 53-57 under 35 U.S.C. §102(b) or 103 has been adequately addressed in view of the amendments to the claims and remarks set forth above. Therefore, the Examiner is respectfully requested to withdraw the rejection and allow the application to proceed to issue.


### III. CONCLUSION

Applicants submit that all of the claims are in condition for allowance, which action is requested. If the Examiner finds that a telephone conference would expedite the prosecution of this application, the Examiner is invited to telephone the undersigned at the number provided.

The Commissioner is hereby authorized to charge any underpayment of fees associated with this communication, including any necessary fees for extensions of time, or credit any overpayment to Deposit Account No. 50-0815, order number GRUE003.

Respectfully submitted,  
BOZICEVIC, FIELD & FRANCIS LLP

Date: December 23, 2003

By:   
Edward J. Baba  
Registration No. 52,581

BOZICEVIC, FIELD & FRANCIS LLP  
200 Middlefield Road, Suite 200  
Menlo Park, CA 94025  
Telephone: (650) 327-3400  
Facsimile: (650) 327-3231



U.S. App. No.: 09/269,874  
Group Art Unit: 1641  
Eaminer: P.A. Duffy  
Title: Recombinant Process for Preparing A Complete Malaria Antigen,  
GP190/MSP1



**Protein:** gp190/MSP1 – amino acids 1-1639 of SEQ ID NO:3  
**Gene:** gp190s  
**Description:** The complete protein with the attachment signal and signal peptide  
**Support:** Specification pages 8 and 15



**Protein:** gp190/MSP1 – amino acids 1-1621 of SEQ ID NO:3  
**Gene:** gp190<sup>s1</sup>  
**Description:** The protein with the signal peptide but lacking an attachment signal  
**Support:** Specification pages 8 and 15



**Protein:** gp190/MSP1 – amino acids 20-1621 of SEQ ID NO:3  
**Gene:** gp190<sup>s2</sup>  
**Description:** The protein lacking the attachment signal and signal peptide  
**Support:** Specification pages 8 and 15

EXPERIMENTAL PARASITOLOGY 67, 1-11 (1988)

Notice: This material may be  
protected by copyright law  
(Title 17 U.S. Code).

*Plasmodium falciparum*: Gene Structure and Hydropathy Profile of  
the Major Merozoite Surface Antigen (gp195) of the Uganda-Palo  
Alto Isolate

SANDRA P. CHANG, KENTON J. KRAMER, KAREN M. YAMAGA, ANN KATO,  
STEPHEN E. CASE, AND WASIM A. SIDDIQUI

Department of Tropical Medicine and Medical Microbiology, University of Hawaii School of Medicine,  
Honolulu, Hawaii, 96816, U.S.A.

(Accepted for publication 19 April 1988)

CHANG, S. P., KRAMER, K. J., YAMAGA, K. M., KATO, A., CASE, S. E., AND SIDDIQUI, W. A. 1988. *Plasmodium falciparum*: Gene structure and hydropathy profile of the major merozoite surface antigen (gp195) of the Uganda-Palo Alto isolate. *Experimental Parasitology* 67, 1-11. The gene encoding the 195,000-Da major merozoite surface antigen (gp195) of the FUP (Uganda-Palo Alto) isolate of *Plasmodium falciparum*, a strain widely used for monkey vaccination experiments, has been cloned and sequenced. The translated amino acid sequence of the FUP gp195 protein is closely related to the sequences of corresponding proteins of the CAMP (Malaysia) and MAD-20 (Papua New Guinea) isolates and more distantly related to those of the Wellcome (West Africa) and KI (Thailand) isolates, supporting the proposed allelic dimorphism of gp195 within the parasite population. The prevalence of dimorphic sequences within the gp195 protein suggests that many gp195 epitopes would be group-specific. Despite the extensive differences in amino acid sequence between gp195 proteins of these two groups, the hydropathy profiles of proteins representative of both groups are very similar. The conservation of overall secondary structure shown by the hydropathy profile comparison indicates that gp195 proteins of the various *P. falciparum* isolates are functionally equivalent. This information on the primary structure of the FUP gp195 protein will enable us to evaluate the possible roles of conserved, group-specific and variable epitopes in immunity to the blood stage of the malaria parasite. © 1988 Academic Press, Inc.

INDEX DESCRIPTORS AND ABBREVIATIONS: *Plasmodium falciparum*; Protozoa, parasitic; Malaria, human; Merozoite; Vaccine; Major merozoite surface antigen (gp195); Uganda-Palo Alto isolate (FUP); Dalton (Da); Kilodalton (kDa); Kilobase (kb); Deoxyribonucleic acid (DNA); Base pair (bp).

# INTRODUCTION

*Plasmodium falciparum* is the causative agent of the most serious form of human malaria. The surface of the malaria parasite undergoes drastic antigenic changes during its complex life cycle. The predominant surface antigen of the sporozoite produced during the sexual cycle of the parasite in the *Anopheles* mosquito host is the circumsporozoite protein (Nussenzweig and Nussenzweig 1985). After injection of the sporozoite into the bloodstream of the vertebrate host and its uptake into hepatocytes the circumsporozoite protein is lost (Dan-

forth *et al.* 1978). Little is known about surface antigens of the parasite during the hepatic stage of asexual development. However, surface antigens of the erythrocytic stages of the parasite life cycle have been well studied. The major surface antigens of the *Plasmodium falciparum* merozoite, the erythrocytic invasive stage of the parasite, are derived from a precursor glycoprotein with a molecular weight of 185-195,000 (gp195) (Freeman and Holder 1983; Hall *et al.* 1983; Holder and Freeman 1984). The gp195 precursor protein is synthesized during the late erythrocytic stage of development and is proteolytically processed to

m or by any  
information  
it owner.

n this journal  
e for personal  
asent is given  
the Copyright  
for copying  
consent does  
m, for adver-  
le. Copy fees  
s on the title

IN 55802  
countries  
countries

iting to the  
the office of  
of address  
oth old and  
parasitology.

lower molecular weight fragments (Freeman and Holder 1983; Hall *et al.* 1984a). Three of these fragments of 83,000, 42,000, and 19,000 Da have been detected on the merozoite surface (Holder and Freeman 1984). An approximately 80,000-Da processing fragment of the gp195 has been localized in the surface coat of the merozoite by immunoelectron microscopy (Heidrich *et al.* 1986).

The genes encoding the gp195 protein of several parasite isolates have been cloned and sequenced, revealing sequence polymorphisms among different genes (Hall *et al.* 1984b; Holder *et al.* 1985; Mackay *et al.* 1985; Weber *et al.* 1986; Tanabe *et al.* 1987; Peterson *et al.* 1988). Gp195-related polypeptides are candidates for a blood stage human malaria vaccine. Results of three monkey vaccination experiments using gp195-derived immunogens showed partial (Hall *et al.* 1984b; Perrin *et al.* 1984) to complete protection (Siddiqui *et al.* 1987). In each experiment, challenge has been with the *P. falciparum* FUP isolate, but only in the latter experiment were the monkeys immunized with gp195 purified from FUP parasites. In order to develop a fully protective recombinant polypeptide or synthetic peptide vaccine based on gp195 and to evaluate the significance of antigenic polymorphism of this protein in protective immunity, it appears crucial to define the structure of the FUP gp195 gene.

This study presents the DNA sequence of the FUP gp195 gene and compares its translated amino acid sequence to others that have been published. In addition, the amino acid sequence dimorphism of gp195 proteins is correlated to secondary structure as predicted by hydropathy analysis.

## MATERIALS AND METHODS

**Parasites.** The Uganda-Palo Alto (FUP) strain of *Plasmodium falciparum* was originally isolated from a patient who had contracted the infection in Uganda and was hospitalized at Stanford Medical Center, Palo

Alto, California, in 1966. In 1967, blood-induced infections with this isolate were established in *Aotus trivirgatus* monkeys at Stanford (Geiman and Meagher 1967). The FUP strain was maintained by serial passage in *Aotus* monkeys by Dr. Schmidt of the Southern Research Institute, Birmingham, Alabama. In 1970, the FUP monkey-passaged strain was obtained from Dr. Schmidt and maintained in this laboratory at the University of Hawaii by serial passage in *Aotus* monkeys. In 1977, continuous *in vitro* cultures of the FUP strain in human erythrocytes were established at the University of Hawaii and have been maintained in this laboratory since that time. The FUP parasites used in this study were derived from *in vitro* cultures.

**Isolation of *P. falciparum* DNA.** DNA was isolated from cultured FUP strain *P. falciparum* using the Trager and Jensen (1976) culture technique with modifications (Siddiqui and Palmer 1981) and standard DNA extraction methods (Maniatis *et al.* 1982). The Protoclone bacteriophage  $\lambda$  gt10 system (Promega Biotec, Madison, WI, U.S.A.) was used to generate an FUP *P. falciparum* genomic library. FUP DNA (0.3  $\mu$ g, 0.1 pmole) was digested with 3 units of the restriction endonuclease *EcoRI* (Boehringer Mannheim, Indianapolis, IN, U.S.A.) and ligated to 0.5  $\mu$ g (0.17 pmole) *EcoRI*-digested  $\lambda$  gt10 DNA (Promega Biotec) with 1 unit T4 polynucleotide ligase (Promega Biotec). The resultant recombinant phage were grown in *Escherichia coli* strain C600AHFL cells, and generated a library of  $2.5 \times 10^6$  plaque-forming units in which 87% of the phage contained inserts.

**Preparation of synthetic oligonucleotides.** Oligonucleotides used as hybridization probes were synthesized as pairs of 30-mers overlapping by 10 base pairs. Probes were synthesized using methoxy phosphoramidites or  $\beta$ -cyanoethyl phosphoramidites on Applied Biosystems DNA synthesizers (Foster City, CA, U.S.A.) according to the manufacturers recommendations. The oligonucleotides were radiolabeled by fill-in reactions using high specific activity (3000 Ci/mmole) 5'-[ $\alpha$ - $^{32}$ P]deoxycytidine triphosphates and -deoxyadenosine triphosphates (Amersham, Arlington Heights, IL, U.S.A.) and Klenow DNA polymerase I (Boehringer Mannheim). Hybridization to phage DNA on nitrocellulose filters and washing were carried out as described by Ullrich *et al.* (1984).

**Restriction map analysis and DNA sequencing.**  $\lambda$  gt10 recombinant phage inserts were subcloned into pUC plasmids for restriction mapping and into M13mp18 or M13mp19 phage for DNA sequencing. M13 subclones were sequenced by the enzymatic method (Sanger *et al.* 1977) using a universal M13 primer (Pharmacia, Piscataway, NJ, U.S.A.) or specific primers complementary to the insert sequence. Sequencing primers were synthesized using  $\beta$ -cyanoethyl phosphoramidites as described above for oligonucleotide probes. DNA sequence data were analyzed

using computer resources sponsored BIONET National Molecular Biology. Proteinally analyzed using the con by Pauletti *et al.* (1985) and

## RESULTS

A  $\lambda$  gt10 library of DNA was screened with oligonucleotide probe served sequences in the Wellcome and K1 strains (1985; Mackay *et al.* 1985; Mackay *et al.* 1985) (10-1, 3-1, and 18-1) coding region of the *gp195* gene (Fig. 1) and were mapped with endonucleases and subcloning vectors. These three  $\lambda$  clones were

The complete DNA acid translation of the gene is presented in Figure 2. The molecular weight of the protein is 196,245. There are glycosylation sites (Sn) in the protein. The protein contains 20 cysteines are conserved in the protein and 13 are located in the hydrophobic carboxyl




FIG. 1. Map of the gp195 gene. The shaded box shows the positions of the ATG start codon. The map indicates the restriction sites of the 18-1 and 10-1 clones. The arrows indicate the sequencing primers. Restriction endonuclease sites are indicated by arrows.

*P. falciparum* MAJOR MEROZOITE SURFACE ANTIGEN

3

blood-induced infected in *Aotus tri-*  
*eiman* and Meagher  
 gained by serial pas-  
 smid of the Southern  
 Alabama. In 1970,  
 was obtained from  
 is laboratory at the  
 sage in *Aotus mon-*  
 cultures of the FUP  
 e established at the  
 in maintained in this  
 P parasites used in  
 ro cultures.

DNA was isolated  
*iparum* using the  
 chnique with mod-  
 1981) and standard  
*s et al.* 1982). The  
 item (Promega Bio-  
 ted to generate an  
 y. FUP DNA (0.3  
 units of the restric-  
 er Mannheim, In-  
 d to 0.5 µg (0.17  
 (Promega Biotec)  
 (Promega Biotec).  
 re grown in *Esch-*  
 and generated a  
 nts in which 87%.

*otides*. Oligonu-  
 bes were synthe-  
 by 10 base pairs.  
 oxy phosphora-  
 dities on Applied  
 oster City, CA,  
 ers recommenda-  
 labeled by fill-in  
 (3000 Ci/mmol)  
 es and -deoxy-  
 am, Arlington  
 A polymerase I  
 n to phage DNA  
 were carried out

sequencing. λ  
 subcloned into  
 ping and into  
 A sequencing.  
 the enzymatic  
 universal M13  
 J.S.A.) or spe-  
 sert sequence.  
 using β-cyano-  
 above for oligo-  
 were analyzed

using computer resources provided by the N.I.H.  
 sponsored BIONET National Computer Resource for  
 Molecular Biology. Protein sequences were addition-  
 ally analyzed using the computer programs described  
 by Pauletti *et al.* (1985) and Gotoh (1986).

## RESULTS

A λ gt10 library of FUP strain genomic  
 DNA was screened with several synthetic  
 oligonucleotide probes based on the con-  
 served sequences in the gp195 genes of the  
 Wellcome and K1 strains (Holder *et al.*  
 1985; Mackay *et al.* 1985). Three λ clones  
 (10-1, 3-1, and 18-1) contained the entire  
 coding region of the strain FUP gp195 gene  
 (Fig. 1) and were mapped using restriction  
 endonucleases and subcloned into M13 se-  
 quencing vectors. The M13 subclones of  
 these three λ clones were sequenced.

The complete DNA sequence and amino  
 acid translation of the strain FUP gp195  
 gene is presented in Fig. 2. The calculated  
 molecular weight of the entire protein is  
 196,245. There are 15 potential N-  
 glycosylation sites (Snider 1984). The pro-  
 tein contains 20 cysteine residues; 19 cys-  
 teines are conserved among various iso-  
 lates and 13 are located immediately before  
 the hydrophobic carboxy terminal region.

The amino acid translation of the FUP  
 gp195 gene has been aligned with the se-  
 quences of other isolates (Fig. 3) using the  
 computer algorithm described by Gotoh  
 (1986). Based on this alignment the gp195  
 sequence can be divided into three types of  
 regions. The first type is the conserved re-  
 gion (85–100% sequence identity). Con-  
 served regions are located at the amino ter-  
 minus and the carboxy terminus, as well as  
 at internal segments of the protein. The sec-  
 ond type of region is the variable repeat  
 region which differs greatly among isolates  
 in both sequence and length and is located  
 toward the amino terminus. The third type  
 of region is designated group-specific be-  
 cause it appears to exist in two forms which  
 differ greatly in sequence (42–46% se-  
 quence identity). Over one-half of the  
 gp195 protein consists of group-specific se-  
 quences. Others have previously recog-  
 nized this pattern of polymorphism of the  
 gp195 gene in the parasite population and  
 refer to it as allelic dimorphism (Tanabe *et*  
*al.* 1987). The FUP protein belongs to the  
 same dimorphic group as the gp195 of the  
 Papua New Guinea isolate MAD20, differ-  
 ing primarily at the variable repeat region.




FIG. 1. Map of the gp195 gene of the Uganda-Palo Alto isolate of *P. falciparum*. The upper striped box shows the positions of the start and stop codons and the length of the coding region. The lower map indicates the restriction endonuclease sites of the λ gt10 clones (3-1 and 10-1) and the M13 subclones of the 18-1 λ gt10 clone (18-1a and 18-1b) shown on the bottom of the figure. The arrows indicate the sequencing strategy using either an M13 universal primer or gp195-specific oligonucleotide primers. Restriction endonucleases: E, *EcoRI*; H, *HindIII*; N, *NdeI*; B, *BglII*; and P, *PstI*.

[illegible]

FIG. 2. Nucleotide sequence of the Uganda-Palo Alto (FUP) gp195 gene. The deduced amino acid sequence of the open reading frame is shown below the nucleotide sequence. The potential signal peptidase site is indicated with an arrowhead at position 57. The variable repeat region is underlined beginning at position 190 and the hydrophobic carboxy terminal sequence is overlined beginning at position 5125. Potential *N*-glycosylation sites are denoted by solid circles.

Several substitutions in the FUP amino acid sequence which influence the sequence, alone or in combination, relative to MAF, alter the overall single amino acid sequence throughout the FUP and MAF. The amino acid sequence is very similar overall, but in contrast to the relationship of FUP and MAF. The variable region of gp195 protein is highly conserved in the lacyan CAMP. In fact, the gp195 protein of the CAMP gp195 is identical to the gp195 of the FUP gene.

The extensive conservation of the group-specific residues may raise the possibility that these residues may represent conserved motifs or motifs. Hydropathic secondary structure elements and interactions (Kyte and Doolittle, 1987) have been used to compare the conservation of amino acid residues in the group-specific regions of the hydropathy profile of the dimorphic gp19 of the measles virus (Kyte and Doolittle, 1985) to determine whether the conservation would be predicted (Fig. 4). As a control, the conserved sequence of the hydropathy profile of the measles virus also differs. However, regions of group-specific conservation of similar hydropathic changes in the hydropathicity interactions differ (1000 (FUP) amino acids) correspond to those that are lacking overall conservation.

deduced amino acid  
The potential signal  
region is underlined  
overlined beginning at

The extensive sequence differences of the group-specific regions of gp195 genes raise the possibility that these dimorphs may represent functionally divergent proteins. Hydropathy profiles, which reflect secondary structure as predicted by water interactions (Kyte and Doolittle 1982), have been used to demonstrate functional conservation among distantly related proteins (Simpson *et al.* 1987). We compared the hydropathy profiles of the FUP and Wellcome dimorphic gp195 proteins using an adaptation of the method of Kyte and Doolittle (Kyte and Doolittle 1982; Pauletti *et al.* 1985) to determine whether these proteins would be predicted to differ in structure (Fig. 4). As anticipated, regions of conserved sequence displayed identical hydropathy patterns. The different repeat regions also differed in hydropathy. However, regions containing dimorphic or group-specific sequences maintained very similar hydropathy patterns with only subtle changes in degree of hydrophilicity or hydrophobicity. A few regions where water interactions differed were at positions 900–1000 (FUP) and 1460–1640 (FUP), which correspond to insertions into the FUP gene that are lacking in the Wellcome gene. The overall conservation of the hydropathy pat-

Sequence comparisons between gp195 genes of different isolates indicate that

amino acid sequences of the dimorphic proteins can differ by greater than 50% in certain parts of the gene. Given such extensive sequence dissimilarities, we examined whether structural differences as reflected by hydropathy profiles could be observed. A comparison of hydropathy profiles of the FUP and Wellcome gp195 proteins, which represent the dimorphic groups, indicated that outside of the variable repeat regions the hydropathy patterns of the two proteins were very similar. These results suggest that the overall structure of the gp195 proteins of different *P. falciparum* isolates is conserved. Such structural conservation suggests that the group-specific regions of gp195 proteins share the same function. However, this function may be more dependent on the overall conformation of the protein than on its primary sequence. Alternatively, it is possible that group-specific regions may be involved in a function which can be carried out using two different pathways, such as the interaction of the parasite with different receptors on the host cell, as has been suggested by others (Tanabe *et al.* 1987).

In designing an effective malaria vaccine, it is essential that the level of antigenic polymorphism of candidate antigens be carefully assessed. It is generally recognized that highly variable, isolate-specific sequences such as the variable repeats of gp195 are less attractive for vaccine development than conserved sequences. Several regions of the gp195 genes are highly conserved among all of the sequences that have been studied. These conserved regions are located on either side of the variable repeats at the amino terminal end of the protein, at the carboxy terminal region, and

between the two large group-specific regions (Fig. 3). These conserved regions would be located on two to three distinct processing fragments on the mature merozoite surface (Lyon *et al.* 1986) and may be important in the merozoite invasion process. However, it remains to be shown whether epitopes contained in these regions are more relevant to immunity than those in other, less-conserved regions. Both conserved and group-specific regions of this protein have a generally hydrophilic character and exposure of these regions on the protein surface would allow them to be recognized as antigenic determinants (Hopp and Woods 1981). The extensive amino acid substitutions in group-specific regions of the gp195 protein make it likely that proteins of different groups would be antigenically distinct. While immunity developed against these regions would probably be group-specific, the evidence that there are a limited number (possibly only two forms) of group-specific regions make it feasible to consider including both of these regions in a recombinant vaccine.

Information on the primary structure of gp195 in the FUP strain enables us to re-evaluate vaccination experiments in which monkeys immunized with antigen from another strain of *P. falciparum* were challenged with parasites of the FUP strain. While it must be recognized that these experiments differed in experimental detail, it is still informative to discuss them in light of this new information. Hall *et al.* (1984b) immunized *Saimiri* monkeys with monoclonal antibody-purified p190, the gp195-equivalent of the K1 (Thailand) strain of the parasite, and challenged these animals with the FUP strain. Two of three immunized

Fig. 3. Comparison of the amino acid sequences encoded by the gp195 gene of the FUP-Uganda, MAD20-Papua New Guinea (Tanabe *et al.* 1987), Wellcome-Lagos (Holder *et al.* 1985), and K1-Thailand (Mackay *et al.* 1985) strains of *P. falciparum*. Alignment was done using the Gotoh algorithm (Gotoh 1986). Shared sequences are indicated by blank spaces and gaps are indicated by periods. Conserved, variable repeat, and group-specific regions are indicated as the respectively designated, overlined sequences.

FUP NAD WEL K1	MRKILL
FUP NAD WEL K1	SOTSS SC V WC V ... V
FUP NAD WEL K1	MLTFL S V
FUP NAD WEL K1	RNKT K K
FUP NAD WEL K1	SGTST T SR N
FUP NAD WEL K1	SLAAD A ... A ...
FUP NAD WEL K1	EDWOC LTWFE LTWFE
FUP NAD WEL K1	NE.FG KIT DI KIT DI
FUP NAD WEL K1	KKEDAV NE KOD NE KOD
FUP NAD WEL K1	TRTYV S LD S LD
FUP NAD WEL K1	IYLYL VON V VON V
FUP NAD WEL K1	SIVSL LFEN LFEN
FUP NAD WEL K1	KLDPLF E E
FUP NAD WEL K1	TLSBVS E E
FUP NAD WEL K1	MTVTV D A D A
FUP NAD WEL K1	DNISGS N E E
FUP NAD WEL K1	LFPSYK F S N F S N
FUP NAD WEL K1	FNDLYL I LSV I LSV
FUP NAD WEL K1	FQMLN LARYVK LARYVK
FUP NAD WEL K1	PLFCGII SHV








FIG. 4. Comparison of hydropathy profiles of the gp195 proteins of the FUP (A) and Wellcome (B) isolates. Hydropathy was analyzed as described by Kyte and Doolittle (1982) using the algorithm adapted by Pauletti *et al.* (1985) for use on a personal computer. Hydrophobic areas are shown above and hydrophilic areas are shown below the horizontal line. Conserved, group-specific, and variable repeat regions were based on the comparison of FUP and Wellcome sequences aligned in Fig. 3.

animals were partially infected (5–10% of the total animal reacted). Immunized controls also reacted. Siddiqui *et al.* (Vietnam) strain profile, also obtained from immunized *Aotus* with FUP strain profile, contrast with the control obtained in *Aotus* from purified FUP gp195 parasites of the same strain (1987). The results of FUP heterologous reactions are particularly interesting for a 36-bp deletion in the first 375 amino acid sequence are 99.9% identical (Fig. 3). The repeat region of the FUP strain is made up of also found in [SAQ(SGT)*n*], where *n* is the larger number of repeats (*n*) and overall the FUP repeat region is of the third type of repeat beyond residue 375. Sequence similarity between proteins drops below 50% protection may have been experiment by in epitopes located within the region of the polypeptide may require conserved epitopes located and/or to conform to the expressed by the long FUP polypeptide. The conserved epitope reported by the findings (1986), in which sequences of gp195 were detected from chlamydia. Most recently have shown that a response to a model

FIG. 4. Comparison of hydropathy profiles of the gp195 proteins of the FUP (A) and Wellcome (B) isolates. Hydropathy was analyzed as described by Kyte and Doolittle (1982) using the algorithm adapted by Pauletti *et al.* (1985) for use on a personal computer. Hydrophobic areas are shown above and hydrophilic areas are shown below the horizontal line. Conserved, group-specific, and variable repeat regions were based on the comparison of FUP and Wellcome sequences aligned in Fig. 3.

animals were partially protected from parasite infection (5–10% parasitemias) while a third animal reacted similarly to the unimmunized controls and required drug treatment. Siddiqui *et al.* (1978), using FVO (Vietnam) strain parasites for immunization, also obtained only partial protection of immunized *Aotus* monkeys challenged with FUP strain parasites. These results contrast with the complete protection obtained in *Aotus* monkeys immunized with purified FUP gp195 and challenged with parasites of the same strain (Siddiqui *et al.* 1987). The results of the strain K1/strain FUP heterologous vaccination experiment are particularly interesting because except for a 36-bp deletion in the repeat region, the first 375 amino acids of K1 and FUP gp195 are 99.9% identical (differ by 16 base pairs, Fig. 3). The repeat region of the challenge FUP strain is made up of repeats which are also found in the K1 polypeptide [SAQ(SGT)<sub>n</sub>], the only differences being the larger number of consecutive SGT repeats (n) and overall greater length of the FUP repeat region and the absence of a third type of repeat unit (SGP). However, beyond residue 375 of the K1 sequence, the sequence similarity between K1 and FUP proteins drops below 50%. While partial protection may have been achieved in this experiment by immunity to conserved epitopes located within the amino terminal region of the polypeptide, complete protection may require immunity to nonconserved epitopes located beyond this region and/or to conformational epitopes expressed by the longer repeat region of the FUP polypeptide. The involvement of nonconserved epitopes in immunity is supported by the findings of Cheung *et al.* (1986), in which *Saimiri* monkeys immunized with a conserved amino terminal peptide of gp195 were also incompletely protected from challenge with the malaria parasite. Most recently Patarroyo *et al.* (1987) have shown that a synthetic peptide corresponding to a moderately conserved region

at the amino terminal end of gp195 (residues 43–53) contributed to the development of protective immunity but could not alone protect *Aotus* monkeys against malaria.

The mounting evidence linking the gp195 molecule to protection in the monkey model (Hall *et al.* 1984b; Perrin *et al.* 1984; Cheung *et al.* 1986; Patarroyo *et al.* 1987; Siddiqui *et al.* 1987), along with the growing number of characterized gp195 genes of different *P. falciparum* isolates, provides a powerful basis for development of a blood stage malaria vaccine. Information on the structural relatedness of the different gp195 genes permits a rational design of vaccination experiments which simulate the polymorphism encountered in nature. It also allows us to evaluate the possible need for a multivalent gp195 vaccine to achieve clinical immunity in a susceptible population.

#### ACKNOWLEDGMENTS

We are grateful to J. Baudler and A. Dasch and Drs. S. Horvath and L. Hood for providing oligonucleotide probes. We thank Drs. L. Tam, G. Hui, L. Simpson, and M. Rittenberg for critical reading of the manuscript. We acknowledge the technical advice of R. Lau and Dr. T. Humphreys during the early stages of this work. This research was supported by United States Agency for International Development Grant DPE-0453-A-00-4039 to W.A.S., University of Hawaii Research, and Training Revolving Fund Grant 21R87860F728B210 to S.P.C., and Bionet Grant 1 U41 RR-01685-03.

#### REFERENCES

- CHEUNG, A., LEBAN, J., SHAW, A. R., MERKLI, B., STOCKER, J., CHIZZOLINI, C., SANDER, C., AND PERRIN, L. H. 1986. Immunization with synthetic peptides of a *Plasmodium falciparum* surface antigen induces antimerozoite antibodies. *Proceedings of the National Academy of Science USA* 83, 8328–8332.
- DANFORTH, H. D., ORIH, A. U., AND NUSSEN-ZWEIG, R. S. 1978. Immunofluorescent staining of exoerythrocytic schizonts of *Plasmodium berghei* in fixed liver tissue with stage-specific immune serum. *Journal of Parasitology* 64, 1123–1125.
- DEGOWIN, R. L. AND POWELL, R. D. 1965. Drug resistance of a strain of *Plasmodium falciparum* from Malaya. *American Journal of Tropical Medicine and Hygiene* 14, 519–528.

- FREEMAN, R. R., AND HOLDER, A. A. 1983. Surface antigens of malaria merozoites. A high molecular weight precursor is processed to an 83,000 mol. wt. form expressed on the surface of *Plasmodium falciparum* merozoites. *Journal of Experimental Medicine* 158, 1647-1653.
- GEIMAN, Q. M. AND MEAGHER, M. 1967. Susceptibility of a New World monkey to *Plasmodium falciparum* from man. *Nature* 215, 437-439.
- GOTOH, O. 1986. Alignment of three biological sequences with an efficient traceback procedure. *Journal of Theoretical Biology* 121, 327-337.
- HALL, R., MCBRIDE, J., MORGAN, G., TAIT, A., ZOLG, J. W., WALLIKER, D., AND SCAIFE, J. 1983. Antigens of the erythrocytic stages of the human malaria parasite *Plasmodium falciparum* detected by monoclonal antibodies. *Molecular and Biochemical Parasitology* 7, 247-265.
- HALL, R., OSLAND, A., HYDE, J. E., SIMMONS, D. L., HOPE, I. A., AND SCAIFE, J. G. 1984a. Processing, polymorphism, and biological significance of P190, a major surface antigen of the erythrocytic forms of *Plasmodium falciparum*. *Molecular and Biochemical Parasitology* 11, 61-80.
- HALL, R., HYDE, J. E., GOMAN, M., SIMMONS, D. L., HOPE, I. A., MACKAY, M., SCAIFE, J., MERKLI, B., RICHLE, R., AND STOCKER, J. 1984b. Major surface antigen gene of a human malaria parasite cloned and expressed in bacteria. *Nature (London)* 311, 379-382.
- HEIDRICH, H.-G., MATZNER, M., MIETTINEN-BAUMANN, A., AND STRYCH, W. 1986. Immunoelectron microscopy shows that the 80,000-dalton antigen of *Plasmodium falciparum* merozoites is localized in the surface coat. *Zeitschrift für Parasitenkunde* 72, 681-683.
- HOLDER, A. A., AND FREEMAN, R. R. 1984. The three major surface antigens on the surface of *Plasmodium falciparum* merozoites are derived from a single high molecular weight precursor. *Journal of Experimental Medicine* 160, 624-629.
- HOLDER, A. A., LOCKYER, M. J., ODINK, K. G., SANDHU, J. S., RIVEROS-MORENO, V., NICHOLLS, S. C., HILLMAN, Y., DAVEY, L. S., TIZARD, M. L. V., SCHWARZ, R. T., AND FREEMAN, R. R. 1985. Primary structure of the precursor to the three major surface antigens of *Plasmodium falciparum* merozoites. *Nature (London)* 317, 270-273.
- HOPP, T. P., AND WOODS, K. R. 1981. Prediction of protein antigenic determinants from amino acid sequences. *Proceedings of the National Academy of Science USA* 78, 3824-3828.
- HUI, G. S. N., AND SIDDIQUI, W. A. 1987. Serum from Pf195 protected *Aotus* monkeys inhibit *Plasmodium falciparum* growth in vitro. *Experimental Parasitology* 64, 519-522.
- KYTE, J., AND DOOLITTLE, R. 1982. A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology* 157, 105-132.
- LYON, J. A., GELLER, R. A., HAYNES, J. D., CHULAY, J. D., AND WEBER, J. L. 1986. Epitope map and processing scheme for the 195,000 dalton surface glycoprotein of *Plasmodium falciparum* merozoites deduced from cloned overlapping segments of the gene. *Proceedings of the National Academy of Science USA* 83, 2989-2993.
- MACKAY, M., GOMAN, M., BONE, N., HYDE, J. E., SCAIFE, J., CERTA, U., STUNNENBERG, H., AND BUJARD, H. 1985. Polymorphism of the precursor for the major surface antigens of *Plasmodium falciparum* merozoites studies at the genetic level. *EMBO Journal* 4, 3823-3829.
- MANIATIS, T., FRITSCH, E. F., AND SAMBROOK, J. 1982. "Molecular Cloning. A Laboratory Manual," pp. 280-281. Cold Spring Harbor Laboratory, New York.
- NUSSENZWEIG, V., AND NUSSENZWEIG, R. S. 1985. Circumsporozoite proteins of malaria parasites. *Cell* 42, 401-403.
- PATARROYO, M. A., ROMERO, P., TORRES, M. L., CLAVIJO, P., MORENO, A., MARTINEZ, A., RODRIGUEZ, R., GUZMAN, F., AND CABEZAS, E. 1987. Induction of protective immunity against experimental infection with malaria using synthetic peptides. *Nature (London)* 328, 629-632.
- PAULETTI, D., SIMMONDS, R., DREESMAN, G. R., AND KENNEDY, R. C. 1985. Application of a modified computer algorithm in determining potential antigenic determinants associated with the AIDS virus glycoprotein. *Analytical Biochemistry* 151, 541-546.
- PERRIN, L. H., MERKLI, B., LOCHE, M., CHIZZOLINI, C., SMART, J., AND RICHLE, R. 1984. Antimalarial immunity in Saimiri monkeys. Immunization with surface components of asexual blood stages. *Journal of Experimental Medicine* 160, 441-451.
- PETERSON, M. G., COPPEL, R. L., MCINTYRE, P., LANGFORD, C. J., WOODROW, G., BROWN, G. V., ANDERS, R. F., AND KEMP, D. J. 1988. Variation in the precursor to the major merozoite surface antigens of *Plasmodium falciparum*. *Molecular and Biochemical Parasitology* 27, 291-302.
- SANGER, F., NICKLEN, S., AND COULSON, A. R. 1977. DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Science USA* 74, 5463-5467.
- SIDDIQUI, W. A., AND PALMER, K. L. 1981. Propagation of malaria parasites in vitro. In "Advances in Cell Culture" (K. Maramorosch, Ed.), Vol. 1, pp. 182-212. Academic Press, New York.
- SIDDIQUI, W. A., SCHNELL, J. V., AND GEIMAN, Q. M. 1972. A model in vitro system to test the susceptibility of human malarial parasites to antimalarial drugs. *American Journal of Tropical Medicine and Hygiene* 21, 392-399.
- SIDDIQUI, W. A., G. S. N., S. P., CHAI, ite surface tects *Aotus parum* malar, emy of Sci. SIDDIQUI, V. KRAMER, F. tial prote vaccinated heterolog Medicine a SIMPSON, L., SIMPSON, AND STUAI cle (mitoch lae and Tr otide sequ 262, 6182-4

*P. falciparum* MAJOR MEROZOITE SURFACE ANTIGEN

11

- ic character of a pro-  
logy 157, 105-132.
- HAYNES, J. D., CHU-  
L. 1986. Epitope map  
195,000 dalton sur-  
fium *falciparum* mero-  
overlapping segments of  
National Academy of
- NE, N., HYDE, J. E.,  
NNENBERG, H., AND  
ism of the precursor  
of *Plasmodium falciparum*  
the genetic level.
- AND SAMBROOK, J.  
Laboratory Manual,"  
por Laboratory, New
- NZWEIG, R. S. 1985.  
malaria parasites. *Cell*
- P., TORRES, M. L.,  
LARTINEZ, A., ROD-  
D CABEZAS, E. 1987.  
munity against experi-  
encing synthetic pep-  
9-632.
- DREESMAN, G. R.,  
pplication of a modi-  
fying potential an-  
with the AIDS virus  
unistry 151, 541-546.
- IE, M., CHIZZOLINI,  
1984. Antimalarial  
Immunization with  
blood stages. *Jour-*  
60, 441-451.
- L., MCINTYRE, P.,  
G., BROWN, G. V.,  
J. 1988. Variation in  
ozoite surface anti-  
*Molecular and Bio-*  
302.
- FULSON, A. R. 1977.  
minating inhibitors.  
*Academy of Science*
- L. 1981. Propaga-  
p. In "Advances in  
1, Ed.), Vol. 1, pp.  
York.
- V., AND GEIMAN,  
tem to test the sus-  
asites to antimalar-  
*Tropical Medicine*
- SIDDIQUI, W. A., TAM, L. Q., KRAMER, K. J., HUI,  
G. S. N., CASE, S. E., YAMAGA, K. M., CHANG,  
S. P., CHAN, E. B., AND KAN, S.-C. 1987. Merozo-  
ite surface coat precursor protein completely pro-  
tects *Aotus* monkeys against *Plasmodium falciparum*  
malaria. *Proceedings of the National Acad-*  
*emy of Science USA* 84, 3014-3018.
- SIDDIQUI, W. A., TAYLOR, D. W., KAN, S.-C.,  
KRAMER, K., AND RICHMOND-CRUM, S. 1978. Par-  
tial protection of *Plasmodium falciparum*-  
vaccinated *Aotus trivirgatus* against a challenge of a  
heterologous strain. *American Journal of Tropical*  
*Medicine and Hygiene* 27, 1277-1278.
- SIMPSON, L., NECKELMANN, N., DE LA CRUZ, B. F.,  
SIMPSON, A. M., FEAGIN, J. E., JASMER, D. P.,  
AND STUART, K. 1987. Comparison of the maxicir-  
cle (mitochondrial) genomes of *Leishmania tarento-*  
*lae* and *Trypanosoma brucei* at the level of nucle-  
otide sequence. *Journal of Biological Chemistry*  
262, 6182-6196.
- SNIDER, M. D. 1984. Biosynthesis of glycoproteins:  
Formation of N-linked oligosaccharides. In "Bi-  
ology of Carbohydrates" (V. Ginsburg and P. Rob-  
bins, Eds.), Vol. 2, pp. 163-198. Wiley, New York.
- TANABE, K., MACKAY, M., GOMAN, M., AND SCAIFE,  
J. G. 1987. Allelic dimorphism in a surface antigen  
gene of the malaria parasite *Plasmodium falciparum*. *Journal of Molecular Biology* 195, 273-287.
- TRAGER, W., AND JENSEN, J. B. 1976. Human malaria  
parasites in continuous culture. *Science* 193, 673-  
675.
- ULLRICH, A., BERMAN, C. H., DULL, T. J., GRAY,  
A., AND LEE, J. M. 1984. Isolation of the human  
insulin-like growth factor gene using a single syn-  
thetic DNA probe. *EMBO Journal* 3, 361-364.
- WEBER, J. L., LEININGER, W. M., AND LYON, J. A.  
1986. Variation in the gene encoding a major mero-  
zoite surface antigen of the human malaria parasite  
*Plasmodium falciparum*. *Nucleic Acids Research*  
14, 3311-3323.

*Molecular and Biochemical Parasitology*, 36 (1989) 271–282  
Elsevier

MOLBIO 01210

Notice: This material may be  
protected by copyright law  
(Title 17 U.S. Code).

271

## Cloning and analysis of the gene encoding the 230-kilodalton merozoite surface antigen of *Plasmodium yoelii*

Alan P. Lewis

Department of Molecular Biology, Wellcome Biotech, Beckenham, Kent, U.K.

(Received 31 March 1989; accepted 16 May 1989)

The complete nucleotide sequence of the gene for the 230-kDa precursor to the major merozoite surface antigens (PMMSA) of *Plasmodium yoelii* YM has been determined. A single open reading frame of 5316 bp encodes a protein of calculated molecular mass 197 000. The deduced amino acid sequence contains potential signal peptide and membrane anchor sequences of 19 and 18 residues, respectively, and a region of six tandemly repeated tetrapeptides, Gly-Ala-Val-Pro. Comparison with the 195-kDa *Plasmodium falciparum* analogue (Pf195) at the amino acid level indicated an overall homology of approximately 30%, with areas of as high as 60% conservation. The tripeptide repeats present near the N-terminus of the Pf195 antigen are absent from the Py230 sequence. The PMMSA can be divided into 22 blocks based upon interspecies amino acid conservation.

Key words: *Plasmodium yoelii*; Malaria; Precursor to the major merozoite surface antigens; Nucleic acid sequence

### Introduction

The precursor to the major merozoite surface antigens (PMMSA) has been proposed as a candidate for a vaccine directed against the asexual erythrocytic stage of malaria [1]. This polypeptide has been identified in human [2,3], simian [4], and rodent [5–7] malarial species, ranging in molecular mass from 185 to 250 kDa. The precursor is synthesised during intraerythrocytic development of the parasite, and is processed into a number of discrete fragments during merozoite maturation [2,8–11]. PMMSA fragments have been detected on the surface coat of the merozoite [12–15] and a possible role for the protein

has been suggested in erythrocyte invasion [16]. The PMMSA has a size of 230 kDa in the murine malaria *Plasmodium yoelii* (Py230) [5]. Purified Py230, and a monoclonal antibody which recognises a C-terminal epitope on this protein, have both been shown to protect mice from challenge infection with *P. yoelii* [5,17,18]. A role for cell-mediated immunity has also been implicated in the response observed to the whole antigen [19]. The analogous protein in the human malaria *Plasmodium falciparum* has a size in the range 185–205 kDa (Pf195) [2]. The complete protein [20–22], or synthetic peptides derived from this antigen [23,24], have been used to produce partial or complete protection against challenge infection in non-human vaccine trials. Furthermore, a polymeric synthetic hybrid protein, based upon a mixture of three synthetic peptides including a derivative of Pf195, has been found to induce protective immunity in humans [25]. These results reinforce the potential of the PMMSA as a candidate for a vaccine against the malarial asexual blood stage, and emphasize the need to develop an experimental model system for this antigen in order to analyse in more detail the mechanisms involved in protective immunity.

Correspondence address: Alan P. Lewis, Dept. of Molecular Biology, Wellcome Biotech, Langley Court, Beckenham, Kent, BR3 3BS, U.K.

Note: Nucleotide sequence data reported in this paper have been submitted to the GenBank™ Data Bank with the accession number J04668.

Abbreviations: PMMSA, precursor to the major merozoite surface antigens; Py230, 230-kDa merozoite antigen of *P. yoelii*; Pf195, 195-kDa merozoite antigen of *P. falciparum*.

0166-6851/89/\$03.50 © 1989 Elsevier Science Publishers B.V. (Biomedical Division)

Krakow, J.L., Eng-  
7) A glycan-phosphatidyl  
D in human serum.

98) A phospholipase  
d anchor of cell-sur-  
roc. Natl. Acad. Sci.

rt, G.W. and Eng-  
or glycan assembly:  
tylinositol anchor of  
coprotein. Cell 56.

T. and Hart, G.W.  
ace glycoprotein of  
63, 17697–17705.

The 3' portion of the gene for the *P. yoelii* 17XL 230-kDa protein was recently published [26]. This study describes the cloning and sequencing of the complete Py230 gene from the YM strain of *P. yoelii*, and a comparison of the sequence with the *P. falciparum* analogue, Pf195.

## Materials and Methods

**Preparation of parasite genomic DNA.** CD1 mice were infected with the YM strain of *P. yoelii*, and lymphocytes removed on day 1 post-infection by subcutaneous injection with 150  $\mu$ l of 40 mg ml<sup>-1</sup> cyclophosphamide in 0.85% NaCl [27]. Parasitised blood was collected on day 3 post-infection, when parasitaemias averaged 50%, and parasite DNA was prepared as previously described [28,29].

**Construction of genomic DNA libraries.** Three libraries were constructed from the *P. yoelii* genomic DNA. All restriction endonuclease digests were under conditions recommended by the manufacturer. Library (1): parasite DNA was digested with mung bean nuclease (Pharmacia) in 35% formamide as described [30], and ligated to phosphorylated *EcoRI* linkers (Pharmacia). After digestion with *EcoRI* (Amersham), excess linkers were removed using a NACS PREPAC column (BRL). The DNA was ligated into the *EcoRI* site of  $\lambda$ gt11 (Stratagene), and a genomic library constructed by in vitro packaging (Stratagene). Library (2): parasite DNA was digested with *DraI* (NBL) and ligated into the *SmaI* site of pUC9 (Pharmacia) treated with calf intestinal alkaline phosphatase [28]. A genomic library was constructed by transformation of Max Efficiency DH5 $\alpha$  competent cells (BRL). Library (3): parasite DNA was digested with *EcoRI* and a genomic library constructed, as above, using *EcoRI*-cut pUC9.

**Screening of genomic libraries.** Libraries were screened using synthetic oligonucleotides made on a Biosearch Sam One DNA synthesiser (New Brunswick). Library (1) was screened using probes A (a 26-mer, 5'-GAAGGTAATACATGTGTAGAAAATAA-3' corresponding to nucleotides 1807-1832 in ref. 26) and B (a 26-mer,

5'-TTTCTTTAACAAGAGAAGAGAA-GCTG-3' corresponding to nucleotides 285-310 in ref. 26); library (2) was screened using probe B, and library (3) using probe C (an 18-mer, 5'-AAACAAAGATGCITTAAG-3' corresponding to nucleotides 2685-2702 in Fig. 2). Bacteriophage plaques and bacterial colonies were lifted on to Hybond-N nylon filters (Amersham) following the manufacturer's instructions, and screened with <sup>32</sup>P-labelled oligonucleotides [31] as previously described [32]. Bacteriophage  $\lambda$  and pUC9 plasmid DNA was isolated from positive clones as described [28,33].

**DNA sequencing.** The insert of the  $\lambda$ gt11 recombinant was subcloned into the *EcoRI* site of pUC9. Regions of the genomic DNA pUC9 clones were sequenced in both directions by plasmid priming following the dideoxy chain termination method [34], according to the Sequenase kit (USB) protocol. Sequences were analysed on the Wellcome Biotech computer system with the aid of the programs IALIGN (National Biomedical Research Foundation) and DIAGON [35].

## Results

**Isolation of *P. yoelii* YM Py230 clones.** The 3' portion of the gene for the Py230 antigen from *P. yoelii* 17XL was published recently [26]. Two oligonucleotide probes, A and B, were synthesised, corresponding to nucleotides 1807-1832 and 285-310, respectively, from the published sequence. These regions were chosen as they possessed high nucleic acid homology to the corresponding sections of the Wellcome Pf195 sequence [36]. Approximately 7  $\times$  10<sup>4</sup> phage from library (1) were screened with probe A, and 10 positive clones were detected. After additional rounds of screening with probes A and B, one recombinant,  $\lambda$ PyM4.3, remained positive.  $\lambda$ PyM4.3 contained an *EcoRI* insert of 4.3 kb (Fig. 1).

Probe B was used to screen approximately 4  $\times$  10<sup>4</sup> recombinants of library (2). One positive clone was isolated, pPyD1.7, which was found to possess a *DraI* insert of 1.7 kb (Fig. 1). From sequence analysis, oligonucleotide probe C was synthesised and used to screen library (3). Approximately 2  $\times$  10<sup>4</sup> recombinants were probed.

Fig. 1. 1  
stop cod  
enzyme s

and 20  
pPyE5.  
*EcoRI*  
A fra  
an RNA  
kb RNA  
the size  
kDa pro  
ing sequ

**Nucleoti**  
gene. T.  
*EcoRI*-c  
mined fr  
the thre  
cleotides  
nucleoti  
ing frame  
codon at  
high, wit  
region ar  
quences.  
the entire  
ing fram  
acid resi  
thus sma  
sodium d  
trophore:  
[36,38-4]  
ancies, a  
petitive r  
amino ac  
of the tel

3AAs-  
des 285-310  
using probe  
18-mer, 5'-  
correspond-  
2). Bacteri-  
s were lifted  
ersham) fol-  
ctions, and  
tides [31] as  
hage  $\lambda$  and  
om positive

gt11 recom-  
ori site of  
DNA pUC9  
ons by plas-  
tain termin-  
sequenase kit  
lysed on the  
with the aid  
Biomedical  
I [35].

nes. The 3'  
gen from *P.*  
6]. Two oli-  
synthesised.  
1-1832 and  
ublished se-  
as they pos-  
o the corre-  
95 sequence  
n library (1)  
10 positive  
al rounds of  
ie recomb-  
yM4.3 con-  
p. 1).  
proximately  
one positive  
as found to  
l). From se-  
obe C was  
ry (3). Ap-  
re probed,

and 20 positives isolated. A genomic DNA clone, pPyE5.9, that was purified, contained a 5.9-kb *EcoRI* insert (Fig. 1).

A fragment from  $\lambda$ PyM4.3 was used to probe an RNA blot of total *P. yoelii* YM RNA. A 7.3-kb RNA species was detected (data not shown); the size expected for a transcript encoding a 230-kDa protein (including 5' leader and 3' non-coding sequences).

**Nucleotide sequence of the *P. yoelii* YM Py230 gene.** The  $\lambda$ PyM4.3 insert was subcloned into *EcoRI*-cut pUC9. The DNA sequence was determined for a region overlapping the inserts from the three recombinant clones, spanning 5775 nucleotides (Fig. 2). A methionine start codon at nucleotide 190 is followed by a single open reading frame of 5316 bp terminating with the first stop codon at nucleotide 5505. The A+T content is high, with an average of 69% within the coding region and 85% for the 5' and 3' untranslated sequences. This is consistent with levels found for the entire *P. yoelii* genome [37]. The open reading frame encodes a polypeptide of 1772 amino acid residues with a calculated size of 197 kDa, thus smaller than the 230 kDa determined using sodium dodecyl sulphate-polyacrylamide gel electrophoresis. A number of other malarial antigens [36,38-41], however, also exhibit such discrepancies, a property thought to be related to the repetitive regions of these polypeptides. The Py230 amino acid sequence contains six tandem repeats of the tetrapeptide Gly-Ala-Val-Pro, which may

account for a similar discrepancy in this antigen. At the N-terminus of the polypeptide is a putative signal peptide of 19 amino acids, and at the C-terminus a potential 18-amino-acid hydrophobic membrane anchor. The sequence contains 20 cysteine residues, of which 10 are situated within the C-terminal 110 amino acids. Of the remaining 10 cysteines, 8 appear to be positioned as 4 pairs, based upon their linear proximities. There are also 11 potential *N*-glycosylation sites (Asn-X-Ser/Thr, where X can be any amino acid with the probable exclusion of proline [42]) scattered throughout the molecule. The sequence of the C-terminal 2310 nucleotides is identical to that published for the Py230 gene from *P. yoelii* 17XL [26]. The virulent YM and 17XL strains were originally derived from a common ancestor, the uncloned avirulent isolate, 17X [43,44].

**Comparisons between the Py230 and Pf195 sequences.** The *P. yoelii* YM Py230 amino acid sequence was aligned with the Wellcome strain Pf195 sequence [36] by computer analysis (Fig. 3; a revised Wellcome Pf195 sequence was used which has been submitted to the GenBank and EMBL databases). An overall homology of 31% was determined, with particular regions exhibiting as much as 60% conservation. 14 of the 20 cysteines within the Py230 sequence are located at positions similar to those in Pf195, including all 10 cysteines at the C-terminus. None of the *N*-glycosylation sites present in either polypeptide, however, are conserved. This may reflect the ob-





Fig. 1. Physical map of the Py230 gene from the YM strain of *P. yoelii*. The upper map indicates the positions of the start and stop codons, and the length of the coding region. The middle section represents a partial restriction map of the gene. Restriction enzyme sites indicated are: B, *Bgl*II; D, *Dra*I; E, *Eco*RI; H, *Hinc*II; N, *Nde*I; Ps, *Pst*I and Pv, *Pvu*II. The lower map shows the genomic DNA clones used for sequence analysis.



**Fig. 2.** Nucleotide sequence of the peptide and its position in the protein.



Fig. 2. Nucleotide sequence of the Py230 gene, and the deduced amino acid sequence of the open reading frame. Putative signal peptide and membrane anchor sequences are boxed and underlined, respectively, whilst the tetrapeptide repeats are overlined. Positions of cysteine residues are denoted by open circles, and potential *N*-glycosylation sites by open diamonds.

276

1 10 20 30 40 50 60 70  
230 MKVIGLLFSFVFFAIKCKSETTEVYNDLIQKLEKLESLSVDGLELFQKSGVLIINATOPTETIDPFTN---  
195 MKIIFFLCSFLFIINTQCVTHESYQELVKKLEALEDAVLTVGYSLFQKMKVNLNBTSGTAVTTSTPGSK  
71 80 90 100 110 120 130 140  
230 -----HNFAQQVQDFVTKFE  
195 GSVASGGSGGSGVASGGSGVASGGSGVASGGSGNSRRTNPSDSSSDSAKSYADLKHVRVNYLLTIK  
141 150 160 170 180 190 200 210  
230 GLGFTETELVNLKALTPNRYGVKYLIESKEEFNGLMHAINFYDVLDRKLNHCANNYCEIPEHLKIS  
195 ELKYPQLDLTNHMLTCDNIHGFYKLIQGYEINELLYKLNIFYDILLRAKLNDCANDYCGIPFNLIKIR  
211 220 230 240 250 260 270 280  
230 BEETEMLKVVILGYRKPIENIQDDIEKLEIYIERNKETVAALNALIAETKKIQPEGNECDNDASCDSDK  
195 ANELDVLLKLVFGYRKPLONIKDNVGMEDYIKNNKKTIEINELIBESKKTIDKNKN-----ATK  
281 290 300 310 320 330 340 350  
230 YNKKKPIYQAMYNVIFYKKQLAETQKVVEVLEKRVSTLKKNDKPLKQIEVLNAPVVTATQIVTGG  
195 EEEKKKLYQAQYDLSIYNKQLEEAHNLISVLEKRIDTLKKENIKELLDKINEIKNPPANSN-----  
351 360 370 380 390 400 410 420  
230 QSSTEPGSGSSASGTSSSGQASAGTQVEQANTVASVTVTSPVGQNGEASTNPQTAQVQPVPTLTIREKQ  
195 -----TPNTLLDKN  
421 430 440 450 460 470 480 490  
230 KKIAQLYAIKEIAKTIKFNLEGIFVDPIELEYFKKKEKESCNLSTSSCKKNKASETIIPLTIRYPNGI  
195 KKISEKEKEIAKTIKFNIDSLFTDPLELEYLRSKKNKIDISAKVETKESTEPNE-----YPNGV  
491 500 510 520 530 540 550 560  
230 SYPLPENDVYNKIANNAAETTYGDLTHP-DNTPLTGDLATNEQARKOLIKAIKKKIKAEKKLETLTNY  
195 TYPLSYNDINNALNELNS---FGDLINPFDYTKEPSKNIYTDNERKKFINEIKERIKIEKKKIESDKSY  
561 570 580 590 600 610 620 630  
230 DNKLETFNQKTPFKEAAKEFYESKFRNKLTSIEIFEKFKTKRDEYMTKKTENLC---EYNTKELINKLN  
195 EDRSKSLNDITKEYEKLLEIYDSKFNNNIDLTNFEKMMGKRYSYKVEKLTHHNTFASYENSKHNLEKLT  
631 640 650 660 670 680 690 700  
230 KQLNLYQDYSLRKDIISNEIEYFSNKKKELQYNINRLAEAVQAKQNVLVASKD-----VPLSTL  
195 KALKYMEDYSLRNIVGKELKYKLNLSIKIENEIETLVENIKKDEEQLFEKKITKDNKPDEKILEVSDI  
701 710 720 730 740 750 760 770  
230 IVELQIQSLLTKQIBQLNKTEVSLNKAQLKDKLYVPKTYGNEGKPEPYLLIYVKKVDRLAQFIPKDESM  
195 VKVQVQVLLMKNIDELKKTQLILKNVELKHNIHVPNSYKQENKQEPYLLVLKKEIDKLVFMPKVESL  
771 780 790 800 810 820 830 840  
230 IAKEKERHEQGPATGESEEVPSGSAESSTDRSTQSSTSSSSSSSTPAAAESSATLPEAPAPAEAS  
195 INEEKKNIK---TEGQSDNSEPSTEGEITGQATTKPGQAGSALGDSVQAQAEQKQAPPPV  
841 850 860 870 880 890 900 910  
230 PSTEASEETTIPPTTQETQPSQAASSTTPAKPVMTKLYVLEKLQKFLVFSYSCHKYVLLQNSTINKDALS  
195 -----VPVPEAKAQVTPPAPVNNKTENVSKLDYLEKLYEFLNTSYCHKYILVSHSTHNEKILK  
911 920 930 940 950 960 970 980  
230 KYALTSEEDKIRTLKRCSELDVLLAIQNNMPTHYSLYESIYDGLQNIYTELEYKEMHYHYKLDENPSI  
195 QYKITKEES--KLSSCDPLDLENIQNNIPVHYSHFDSLNSLSQLEMEIYEKEMVONLYKLDNDKIK

Fig. 3. Com  
using the [AI  
tween sequen  
sylation sites  
amino acids b




Fig. 3. Comparison of the amino acid sequence of Py230 with Pf195 from the Wellcome strain [36]. Alignment was carried out using the IALIGN program (National Biomedical Research Foundation). Positions of cysteine residues that are conserved between sequences are indicated by filled circles, and those that are not conserved by open circles. Positions of potential N-glycosylation sites are denoted by open diamonds. The positions of previously determined Pf195 blocks based upon conservation of amino acids between different Pf195 alleles [47] are shown. Conserved blocks are boxed by unbroken lines, semi-conserved blocks by broken lines, and variable blocks remain unboxed.

servation that the Py230 antigen is lacking in glycosylation [8], and that the glycosylation identified in the Pf195 may be confined to a glycolipid anchor identified at the C-terminus [45,46]. The Py230 sequence, when compared to Pf195, possesses two large 'inserted' blocks of amino acids in the central region and near to the N-terminus of the polypeptide. There is also a large deletion of residues following the putative signal peptide, which spans the tripeptide repeats (Ser-X-X) present in the Wellcome Pf195.

The Py230 amino acid sequence was compared

to the two Pf195 allelic variants from the Wellcome and MAD20 [47] strains of *P. falciparum*, using the DIAGON computer program of Staden [35] (Fig. 4). Regions of conservation between Py230 and the Wellcome Pf195 allele were found to be similarly conserved when the MAD20 Pf195 allele was compared. The Py230 and Wellcome Pf195 sequences have also been compared to the published sequence for a portion of the 200-kDa PMMSA from the human malaria *Plasmodium vivax* [3] (data not shown). This aligns with the region 117–794 amino acid residues of Py230. It

was found  
between the  
closely mi  
of the oth

## Discussion

The data and sequence analysis of the 195-kDa meroprotein has shown that the 195-kDa meroprotein has a similar putative anchor sequence to the other meroproteins. The level of expression of the 195-kDa meroprotein in Py230 as a control was determined by Western blotting and PMMSA.

Certainly, between the *P. yoelii*, *P. berghei* parasites and other groups, position and malaria genes [situated in a amino acids the three ev- gues that the these region:

The *P. fal* previously been conservation Pf195 alleles ified as either (ogy), semi-co or variable (the alignment may not nec tion of amin




Fig. 4. Comparisons of the Py230, Wellcome Pf195 [36] and MAD20 Pf195 [47] amino acid sequences. Analysis was by the DIAGON program of Staden [35], using a proportional algorithm (proportional score 132, span length 11). The axes of the plots represent the appropriate amino acid sequences numbered from their N- to C-termini.

Fig. 5. Represent  
The sequence is a  
possessing bet

n the Well-  
falciparum,  
m of Staden  
on between  
were found  
AD20 Pf195  
Wellcome  
ared to the  
ie 200-kDa  
lasmodium  
is with the  
(Py230, li

was found that conservation of amino acids between the PMMSAs of any two of the species closely mirrored the homologies observed for each of the other species comparisons.

### Discussion

The data presented here describe the cloning and sequencing of the complete gene for the 230-kDa merozoite antigen from *P. yoelii* YM. The protein has a structure closely resembling that of the 195-kDa PMMSA of *P. falciparum*, with similar putative signal peptide and membrane anchor sequences, and extensive amino acid homologies throughout the length of the molecule. The level of similarity consolidates the use of the Py230 as an experimental model system for the PMMSA.

Certain protein sequences are conserved between the PMMSAs of the three malarial species *P. yoelii*, *P. falciparum* and *P. vivax*: Malaria parasites are thought to fall into three evolutionary groups, based upon genomic DNA base composition and sequence similarities between malaria genes [48,49], and each of the above species situates in a separate group. The conservation of amino acids observed between the PMMSAs of the three evolutionarily distant species thus argues that there are certain constraints placed upon these regions of the polypeptide.

The *P. falciparum* PMMSA sequence has previously been divided into 17 blocks based upon conservation of amino acids between different Pf195 alleles [47]. These regions have been classified as either conserved (more than 87% homology), semi-conserved (areas of patchy homology) or variable (extensive divergence). As shown by the alignment in Fig. 3, however, these divisions may not necessarily reflect the actual conservation of amino acid sequences between malarial

species. Certain 'variable' blocks from the Pf195 allelic analysis, such as amino acids 385-608 from the Wellcome sequence, can be seen to contain regions of close homology when Py230/Pf195 comparisons are made. By contrast, 'conserved' blocks can possess areas of comparatively little homology. DIAGON analysis indicates that any homologies observed between Py230 and Pf195 can also be reproduced in comparisons between the different Pf195 alleles from the Wellcome and MAD20 strains, even within the 'variable' blocks (Fig. 4). This result suggests that the interspecies conservation can be taken as a good indication of the PMMSA regions that are associated with essential structural and/or functional roles.

The Py230 amino acid sequence can be divided into 22 different blocks based upon interspecies conservation (Fig. 5). The blocks are classified thus; (a) conserved (possessing greater than 45% homology); (b) semi-conserved (between 20 and 45% homology); and (c) variable (less than 20% homology and frequently containing large deletions or insertions of amino acids). All of the conserved cysteine residues are found within conserved blocks, thus suggesting important structural functions for these amino acids. The semi-conserved blocks often contain sequences of low homology interspersed with small regions of high conservation, and the latter sequences again probably signify amino acids of physical importance to the protein. The conserved and semi-conserved blocks are essentially  $\alpha$ -helical in structure with the variable regions consisting of randomly coiled hydrophilic amino acids (data not shown). Such data suggest that the conserved blocks represent sequences internal to the protein, with the variable regions positioned on the extremities at the apices of adjacent  $\alpha$ -helices. This would allow for the sequence variability observed. Exceptions to the rule are the repeat re-




Fig. 5. Representation of the Py230 protein sequence based upon amino acid conservation between the Py230 and Pf195 antigens. The sequence is divided into conserved blocks of greater than 45% amino acid homology (black boxes), semi-conserved blocks possessing between 20 and 45% homology (shaded boxes), and variable blocks of less than 20% homology (open boxes).

i [36] and  
as by the  
tional] al-  
ie axes of  
sequences

gions, Gly-Ala-Val-Pro and Ser-X-X, present in variable blocks of Py230 and Pf195, respectively. These are hydrophobic in each case, suggesting that they may be positioned on the interior of the protein. It has been proposed that tandem repeats in malaria antigens are likely to possess some major role, perhaps playing a critical part in the specific functioning of the antigen, or acting as an immunological decoy directing immune responses away from more functionally important regions of the polypeptide [49,50]. The assumed deep-seated positions of these structures, however, combined with the fact that repeats present in the PMMSA of one species are lacking in that of the other species, argues against both of these suggested roles. Indeed, a Pf195 antigen has been described that is totally lacking in repeats [51]. Whether the tandem repeats of the PMMSA possess an important function thus remains unclear.

The external positioning of the variable regions within the PMMSA makes them potentially highly immunogenic. The very variability of such

## References

- Gly-Ala-Val-Pro and Ser-X-X, present in variable blocks of Py230 and Pf195, respectively. These are hydrophobic in each case, suggesting that they may be positioned on the interior of the protein. It has been proposed that tandem repeats in malaria antigens are likely to possess some major role, perhaps playing a critical part in the specific functioning of the antigen, or acting as an immunological decoy directing immune responses away from more functionally important regions of the polypeptide [49,50]. The assumed deep-seated positions of these structures, however, combined with the fact that repeats present in the PMMSA of one species are lacking in that of the other species, argues against both of these suggested roles. Indeed, a Pf195 antigen has been described that is totally lacking in repeats [51]. Whether the tandem repeats of the PMMSA possess an important function thus remains unclear.
- The external positioning of the variable regions within the PMMSA makes them potentially highly immunogenic. The very variability of such
- sequences, however, argues against them as being effective in a malarial vaccine. Two T-cell epitopes have been identified within the N-terminal portion of the Pf195 [52], and both are situated in a conserved region of the PMMSA (Fig. 5; block 3). The conserved blocks could perhaps be used as a basis to limit the search for valuable PMMSA B- and T-cell epitopes, as such epitopes, within these areas, may be expected to show little or no intraspecies antigenic variation.
- ### Acknowledgements
- I would like to thank Steve Nicholls for assistance with computing, Hugh Spence for oligonucleotide synthesis, Dr. Mike Lockyer for critical reading of the manuscript, Dr. Tony Holder for helpful discussions, and Pam Sparks for secretarial help. This work was supported by a United Nations Development Program/World Bank/World Health Organization Special Programme for Research and Training in Tropical Diseases.
- ### References
- Holder, A.A. (1988) The precursor to major merozoite surface antigens: structure and role in immunity. *Prog. Allergy*, 41, 72-97.
  - Holder, A.A. and Freeman, R.R. (1982) Biosynthesis and processing of a *Plasmodium falciparum* schizont antigen recognized by immune serum and a monoclonal antibody. *J. Exp. Med.* 156, 1528-1538.
  - Del Portillo, H.A., Gysin, J., Mattei, D.M., Khouri, E., Udagama, P.V., Mendis, K.N. and David, P.H. (1988) *Plasmodium vivax*: cloning and expression of a major blood-stage surface antigen. *Exp. Parasitol.* 67, 346-353.
  - Epstein, N., Miller, L.H., Kaushel, D.C., Udcinya, I.J., Renner, J., Howard, R.J., Asofsky, R., Aikawa, M. and Hess, R.L. (1981) Monoclonal antibodies against a specific surface determinant on malarial (*Plasmodium knowlesi*) merozoites block erythrocytic invasion. *J. Immunol.* 127, 212-217.
  - Holder, A.A. and Freeman, R.R. (1981) Immunization against blood-stage rodent malaria using purified parasite antigens. *Nature* 294, 361-364.
  - Boyle, D.B., Newbold, C.I., Smith, C.C. and Brown, K.N. (1982) Monoclonal antibodies that protect in vivo against *Plasmodium chabaudi* recognize a 250 000-dalton parasite polypeptide. *Infect. Immun.* 38, 94-102.
  - Newbold, C.I., Boyle, D.B., Smith, C.C. and Brown, K.N. (1982) Identification of a schizont and species specific surface glycoprotein on erythrocytes infected with rodent malarials. *Mol. Biochem. Parasitol.* 5, 45-54.
  - Holder, A.A. and Freeman, R.R. (1984) Characterisation of a high molecular weight protective antigen of *Plasmodium yoelii*. *Parasitology* 88, 211-219.
  - David, P.H., Hadley, T.J., Aikawa, M. and Miller, L.H. (1984) Processing of a major parasite surface glycoprotein during the ultimate stages of differentiation in *Plasmodium knowlesi*. *Mol. Biochem. Parasitol.* 11, 267-282.
  - Hall, R., Osland, A., Hyde, J.E., Simmons, D.L., Hope, I.A. and Scaife, J. (1984) Processing, polymorphism and biological significance of P190, a major surface antigen of the erythrocytic forms of *Plasmodium falciparum*. *Mol. Biochem. Parasitol.* 11, 61-80.
  - Lyon, J.A., Geller, R.H., Haynes, J.D., Chulay, J.D. and Weber, J.L. (1986) Epitope map and processing scheme for the 195 000-dalton surface glycoprotein of *Plasmodium falciparum* merozoites deduced from cloned overlapping segments of the gene. *Proc. Natl. Acad. Sci. USA* 83, 2989-2993.
  - Atkinson, C.T., Aikawa, M., Fujino, T., Tam, L.Q., Hu, G.S.N. and Siddiqui, W.A. (1987) Ultrastructural localization of protective and non-protective *Plasmodium falciparum* proteins using serum samples from vaccinated Aotus monkeys. *J. Parasitol.* 73, 1235-1240.
  - Howard, R.J., Lyon, J.A., Diggs, C.L., Haynes, J.D., Leech, J.H., Barnwell, J.W., Aley, S.B., Aikawa, M. and Miller, L.H. (1984) Localization of the major *Plasmodium falciparum* glycoprotein on the surface of mature intra-
- erythrocytic  
astol. 11.  
14 Freeman,  
of malar  
is proces  
surface o  
Med. 158  
15 Heidrich,  
and Stryc  
that the f  
merozoite  
72, 681-6  
16 Hadley, 1  
sion of ei  
molecular  
17 Freeman,  
of the pro  
a purified  
Immunol.  
18 Majarian,  
C.A. (198  
with an I  
3131-3137  
19 Playfair, .  
Holder, A  
stage mala  
T cell med  
20 Perrin, L.I  
. J. and Ric  
monkeys.  
sexual bloo  
21 Hall, R., I  
I.A., Mac  
Stocker, J  
malaria pa  
311, 379-3  
22 Siddiqui, V  
Case, S.E  
and Kan,  
protein co  
*modium fe*  
84, 3014-3  
23 Cheung, A  
J., Chizzol  
munizatio  
*ciparum su*  
Proc. Natl  
24 Patarroyo,  
Moreno, A  
Cabezas,  
against exp  
peptides. M  
25 Patarroyo,  
Guzman, I  
rillo, L.A.  
thetic vac  
exual bloo  
Nature 332  
26 Burns Jr., J

## Acknowledgements

I would like to thank Steve Nicholls for assistance with computing, Hugh Spence for oligonucleotide synthesis, Dr. Mike Lockyer for critical reading of the manuscript, Dr. Tony Holder for helpful discussions, and Pam Sparks for secretarial help. This work was supported by a United Nations Development Program/World Bank/World Health Organization Special Programme for Research and Training in Tropical Diseases.

n as being  
T-cell epi-  
terminal  
situated in  
5; block  
s be used  
PMMSA  
es, within  
ttle or no

or assist-  
oligonu-  
critical  
older for  
secretar-  
a United  
m/World  
ial Pro-  
Tropical

cterisation  
of *Plasmo-*

iller, L.H.  
coprotein  
i *Plasmo-*  
7-282.  
L., Hope,  
bism and  
ntigen of  
um Mol.

J.D. and  
g scheme  
smodium  
erlapping  
USA 83,

.Q., Hui,  
al locali-  
ium *fal-*  
ccinated

es: J.D.,  
, M. and  
modium  
: intraer-

- throcytic trophozoites and schizonts. *Mol. Biochem. Parasitol.* 11, 349-362.
- 14 Freeman, R.R. and Holder, A.A. (1983) Surface antigens of malaria merozoites. A high molecular weight precursor is processed to an 83 000-mol.wt form expressed on the surface of *Plasmodium falciparum* merozoites. *J. Exp. Med.* 158, 1647-1653.
  - 15 Heidrich, H.-G., Matzner, M., Miettinen-Baumann, A. and Strych, W. (1986) Immunoelectron microscopy shows that the 80 000-dalton antigen of *Plasmodium falciparum* merozoites is localised in the surface coat. *Z. Parasitenkd.* 72, 681-683.
  - 16 Hadley, T.J., Klotz, F.W. and Miller, L.H. (1986) Invasion of erythrocytes by malaria parasites: a cellular and molecular overview. *Annu. Rev. Microbiol.* 40, 451-477.
  - 17 Freeman, R.R. and Holder, A.A. (1983) Characteristics of the protective response of BALB/c mice immunized with a purified *Plasmodium yoelii* schizont antigen. *Clin. Exp. Immunol.* 54, 609-616.
  - 18 Majarian, W.R., Daly, T.M., Weidanz, W.P. and Long, C.A. (1984) Passive immunization against murine malaria with an IgG3 monoclonal antibody. *J. Immunol.* 132, 3131-3137.
  - 19 Playfair, J.H.L., De Souza, J.B., Freeman, R.R. and Holder, A.A. (1985) Vaccination with a purified blood-stage malaria antigen in mice: correlation of protection with T cell mediated immunity. *Clin. Exp. Immunol.* 62, 19-23.
  - 20 Perrin, L.H., Merkli, B., Loche, M., Chizzolini, C., Smart, J. and Richle, R. (1984) Antimalarial immunity in *Saimiri* monkeys. Immunization with surface components of asexual blood stages. *J. Exp. Med.* 160, 441-451.
  - 21 Hall, R., Hyde, J.E., Goman, M., Simmons, D.L., Hope, I.A., Mackay, M., Scaife, J., Merkli, B., Richle, R. and Stocker, J. (1984) Major surface antigen gene of a human malaria parasite cloned and expressed in bacteria. *Nature* 311, 379-382.
  - 22 Siddiqui, W.A., Tam, L.Q., Kramer, K.J., Hui, G.S.N., Case, S.E., Yamaga, K.M., Chang, S.P., Chan, E.B.T. and Kan, S.-C. (1987) Merozoite surface coat precursor protein completely protects *Aotus* monkeys against *Plasmodium falciparum* malaria. *Proc. Natl. Acad. Sci. USA* 84, 3014-3018.
  - 23 Cheung, A., Leban, J., Shaw, R.A., Merkli, B., Stocker, J., Chizzolini, C., Sander, C. and Perrin, L.H. (1986) Immunization with synthetic peptides of a *Plasmodium falciparum* surface antigen induces antimerozoite antibodies. *Proc. Natl. Acad. Sci. USA* 83, 8328-8332.
  - 24 Patarroyo, M.E., Romero, P., Torres, M.L., Clavijo, P., Moreno, A., Martínez, A., Rodríguez, R., Guzman, F. and Cabezas, E. (1987) Induction of protective immunity against experimental infection with malaria using synthetic peptides. *Nature* 328, 629-632.
  - 25 Patarroyo, M.E., Amador, R., Clavijo, P., Moreno, A., Guzman, F., Romero, P., Tascon, R., Franco, A., Murillo, L.A., Ponton, G. and Trujillo, G. (1988) A synthetic vaccine protects humans against challenge with asexual blood stages of *Plasmodium falciparum* malaria. *Nature* 332, 158-161.
  - 26 Burns Jr., J.M., Daly, T.M., Vaidya, A.B. and Long, C.A. (1988) The 3' portion of the gene for a *Plasmodium yoelii* merozoite surface antigen encodes the epitope recognized by a protective monoclonal antibody. *Proc. Natl. Acad. Sci. USA* 85, 602-606.
  - 27 Berenbaum, M.C. (1975) The clinical pharmacology of immunosuppressive agents. In: *Clinical Aspects of Immunology*, 3rd ed., (Gell, P.G.H., Coombs, R.R.A. and Lachmann, P.J., eds.), pp. 689-758, Blackwell, Oxford, London, Edinburgh and Melbourne.
  - 28 Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982) *Molecular Cloning. A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
  - 29 Martia, W.J., Finerty, J. and Rosenthal, A. (1971) Isolation of *Plasmodium berghei* (Malaria) parasites by ammonium chloride lysis of infected erythrocytes. *Nature New Biol.* 233, 260-261.
  - 30 McCutchan, T.F., Hansen, J.L., Dame, J.B. and Mullins, J.A. (1984) Mung bean nuclease cleaves *Plasmodium* genomic DNA at sites before and after genes. *Science* 225, 625-628.
  - 31 Woods, D. (1984) Oligonucleotide screening of cDNA libraries. *Focus* 6 (3), 1-2.
  - 32 Church, G.M. and Gilbert, W. (1982) Genomic sequencing. *Proc. Natl. Acad. Sci. USA* 81, 1991-1995.
  - 33 Pratt, J.M. (1984) Coupled transcription-translation in prokaryotic cell-free systems. In: *Transcription and translation: a practical approach* (Hames, B.D. and Higgins, S.J., eds.), pp. 179-209, IRL Press, Oxford and Washington, DC.
  - 34 Sanger, F., Nicklen, S. and Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
  - 35 Staden, R. (1982) An interactive graphics program for comparing and aligning nucleic acid and amino acid sequences. *Nucleic Acids Res.* 10, 2951-2961.
  - 36 Holder, A.A., Lockyer, M.J., Odink, K.G., Sandhu, J.S., Riveros-Moreno, V., Nicholls, S.C., Hillman, Y., Davey, L.S., Tizard, M.L.V., Schwarz, R.T. and Freeman, R.R. (1985) Primary structure of the precursor to the three major surface antigens of *Plasmodium falciparum* merozoites. *Nature* 317, 270-273.
  - 37 Chance, M.L., Warhurst, D.C., Baggaley, V.C. and Peters, W. (1972) Preparation and characterisation of DNA from rodent malaras. *Trans. R. Soc. Trop. Med. Hyg.* 66, 3-4.
  - 38 Ozaki, L.S., Svec, P., Nussenzweig, R.S., Nussenzweig, V. and Godson, G.N. (1983) Structure of the *Plasmodium knowlesi* gene coding for the circumsporozoite protein. *Cell* 34, 815-822.
  - 39 Eichinger, D.J., Arnot, D.E., Tam, J.P., Nussenzweig, V. and Enea, V. (1986) Circumsporozoite protein of *Plasmodium berghei*: gene cloning and identification of the immunodominant epitopes. *Mol. Cell. Biol.* 6, 3965-3972.
  - 40 Stahl, H.D., Kemp, D.J., Crewther, P.E., Scanlon, D.B., Woodrow, G., Brown, G.V., Bianco, A.E., Anders, R.F. and Coppel, R.L. (1985) Sequence of a cDNA encoding a small polymorphic histidine- and alanine-rich protein from *Plasmodium falciparum*. *Nucleic Acids Res.* 13, 7837-7846.
  - 41 Favaloro, J.M., Coppel, R.L., Corcoran, L.M., Foote,

- S.J., Brown, S.V., Anders, R.F. and Kemp, D.J. (1986) Structure of the RESA gene of *Plasmodium falciparum*. *Nucleic Acids Res.* 14, 8265-8277.
- 42 Von Heijne, G. (1987) *Sequence Analysis in Molecular Biology: Treasure Trove or Trivial Pursuit*. Academic Press, London.
- 43 Yoeli, M., Hargreaves, B.J., Carter, R. and Walliker, D. (1975) Sudden increase in virulence in a strain of *Plasmodium berghei yoelii*. *Ann. Trop. Med. Parasitol.* 69, 173-178.
- 44 Weinbaum, F.I., Evans, C.B. and Tigelaar, R.E. (1976) An in vitro assay for T cell immunity to malaria in mice. *J. Immunol.* 116, 1280-1283.
- 45 Haldar, K., Ferguson, M.A.J. and Cross, G.A.M. (1985) Acylation of a *Plasmodium falciparum* merozoite surface antigen via sn-1,2-diacyl glycerol. *J. Biol. Chem.* 260, 4969-4974.
- 46 Schwarz, R.T., Lockyer, M.J. and Holder, A.A. (1987) Aspects of the post-translational modification of a major *Plasmodium falciparum* merozoite surface antigen. In: *Host-Parasite Cellular and Molecular Interactions in Protozoal Infections* (Chang, K.P. and Snary, D., eds.), pp. 275-280, Springer-Verlag, Heidelberg.
- 47 Tanabe, K., Mackay, M., Goman, M. and Scaife, J.G. (1987) Allelic dimorphism in a surface antigen gene of the malaria parasite *Plasmodium falciparum*. *J. Mol. Biol.* 195, 273-287.
- 48 McCutchan, T.F., Dame, J.B., Miller, L.H. and Barnwell, J. (1984) Evolutionary relatedness of *Plasmodium* species as determined by the structure of DNA. *Science* 225, 808-811.
- 49 Weber, J.L. (1988) Molecular biology of malaria parasites. *Exp. Parasitol.* 66, 143-170.
- 50 Kemp, D.J., Coppel, R.L. and Anders, R.F. (1987) Repetitive proteins and genes of malaria. *Annu. Rev. Microbiol.* 41, 181-208.
- 51 Certa, U., Rotmann, D., Matile, H. and Reber-Liske, R. (1987) A naturally occurring gene encoding the major surface antigen precursor p190 of *Plasmodium falciparum* lacks tripeptide repeats. *EMBO J.* 6, 4137-4142.
- 52 Crisanti, A., Muller, H.-M., Hilbich, C., Sinigaglia, F., Matile, H., McKay, M., Scaife, J., Beyreuther, K. and Bujard, H. (1988) Epitopes recognized by human T cells map within the conserved part of the GP190 of *P. falciparum*. *Science* 240, 1324-1326.

Molecular  
Elsevier

MOLBIO

Precu

Key words:

The pr  
antigen ( candidate  
*Plasmodi*  
ported the  
gene enc  
rodent m  
the gene  
*yoelii* P19  
tope recc  
tibody [2  
been loca  
antigen (i  
portant e  
we have  
reported  
*P. yoel*  
size-selec  
subgenon  
*NdeI* frag  
insert. Py  
was isolat

Correspond  
and Immun  
Sts., PhiladeNote: Nucl  
been submit  
sion numberAbbreviation  
surface antig

0166-6851/85



*Molecular and Biochemical Parasitology*, 43 (1990) 231–244  
Elsevier

MOLBIO 01416

231

Notice: This material may be  
protected by copyright law  
(Title 17 U.S. Code).

## Molecular cloning and sequence analysis of the gene encoding the major merozoite surface antigen of *Plasmodium chabaudi chabaudi* IP-PC1

Willy Deleersnijder, Diana Hendrix, Najib Bendahman, Josée Hanegreets, Lea Brijs, Cécile Hamers-Casterman and Raymond Hamers

Laboratorium Algemene Biologie, Instituut voor Moleculaire Biologie, Vrije Universiteit Brussel, Brussels, Belgium

(Received 12 March 1990; accepted 27 June 1990)

The complete nucleotide sequence of the gene encoding the precursor to the major merozoite surface antigens of *Plasmodium chabaudi chabaudi* strain IP-PC1 has been determined. A single open reading frame was detected, that coded for a protein of 199 kDa. The encoded protein (p199) contains putative signal and membrane anchor sequences and shows a clustering of Cys residues in the last 120 amino acids. Incompletely conserved tandem repeat oligopeptides are present at different positions in the molecule. P199 shows 69% overall homology to the analogous antigen in *Plasmodium yoelii yoelii* strain YM. The divergence between these antigens is largely confined to 4 areas where a number of insertions and/or deletions have occurred. All repeats occur in these divergent regions. The overall homology with both alleles of *Plasmodium falciparum* PMMSA is 33%.

Key words: Malaria; *Plasmodium chabaudi chabaudi*; Precursor to the major merozoite surface antigens; Nucleic acid sequence; Epitope mapping

### Introduction

At the end of its intraerythrocytic development the *Plasmodium* parasite undergoes several rounds of nuclear division and forms a number of individual merozoites. After rupture of the infected red cell the merozoites are released into the bloodstream and rapidly reinvade new erythrocytes. Since merozoites represent the only stage in the erythrocytic cycle that is directly exposed to the immune system, they are considered to be important targets for vaccination [1].

Correspondence address: Willy Deleersnijder, Laboratorium Algemene Biologie, Instituut voor Moleculaire Biologie, Paardenstraat 65, B-1640 St. Genesius Rode, Belgium

Note: Nucleotide sequence data reported in this paper have been submitted to the GenBank™ data base with the accession number M34947.

Abbreviations: PMMSA, precursor to the major merozoite surface antigens; p199, PMMSA of *P. c. chabaudi* IP-PC1; p197, PMMSA of *P. y. yoelii* YM; mAb, monoclonal antibody; IIF, indirect immunofluorescence; AA, amino acid.

The major surface antigens of the merozoite are derived from a high-molecular-weight precursor glycoprotein, that is often referred to as PMMSA (precursor to the major merozoite surface antigens). This antigen is synthesized late in the erythrocytic cycle and is subsequently processed into a number of smaller protein fragments that are associated with the surface of the mature merozoite [2–7]. The  $M_r$  of the precursor, as determined by SDS-polyacrylamide gel electrophoresis (SDS-PAGE), varies from 185–205 kDa in *Plasmodium falciparum* [8] to 230 kDa (*Plasmodium yoelii yoelii*) [9] or 250 kDa (*P. chabaudi chabaudi*) [10–11]. PMMSA shows considerable size and antigenic polymorphism between different isolates of *P. falciparum* [12] and *P. c. chabaudi* [13]. The PMMSA gene has been cloned and sequenced for a number of different isolates of *P. falciparum* [14–21]. Comparison of these sequences shows that the PMMSA gene consists of blocks that are highly conserved and blocks that vary significantly between isolates [16]. Within each variable block only two distinct sequences have been

Vol. 3, 257–265.  
D.V. (1985) Purification of a bifunctional thymidine kinase from methotrexate-resistant cells. *Journal of Biochemistry* 24, 678–686.  
K. (1984) Purification of a polyoma virus polymerizing enzyme from hamster cells. *Journal of Virology* 48, 505–511.  
owsky, A., and Beveridge, J. (1985) Purification of a methotrexate-resistant *Leishmania* cell line. *Journal of Protozoology* 32, 253–254.  
ann, M.P. (1964) The metabolism of hydroxy-methotrexate. *Journal of Biological Chemistry* 239, 1596–1600.

found with the exception of the variable block closest to the amino-terminus where three versions have been found [22]. Furthermore most sequences contain near the aminoterminal end a region of tripeptide tandem repeats that is highly polymorphic between different strains.

Partial sequence information is also available on the PMMSA gene of *Plasmodium vivax* [23] and the complete sequence of the *P. y. yoelii* YM PMMSA gene has recently been published [24].

Although several research groups have been able to induce partial or complete protection with purified PMMSA [25–28] or synthetic oligopeptides derived from it [29–31], the exact mechanism by which this protective immunity operates is still unclear and probably involves both humoral and cell-mediated immunity [32–35]. As a first step towards developing a mouse model system in which these questions might more easily be addressed we have cloned and sequenced the PMMSA gene of the rodent malaria parasite *P. c. chabaudi* IP-PC1 and also established a crude epitope map of PMMSA.

#### Materials and Methods

**Parasites.** Strain IP-PC1 of *P. c. chabaudi* [36], obtained from Dr. P. Falanga (Institut Pasteur Paris) was cloned by limiting dilution. One clone, termed IP-PC1/C was used for the sequence analysis described in this study. Strain IP-PC1 is a rat-adapted strain that was transferred to mice where it induces fairly synchronous infections. IP-PC1 schizont infected erythrocytes do not sequester. Several attempts to mosquito-transmit IP-PC1 or IP-PC1/C were unsuccessful. Parasites were grown in OF1 outbred mice (Iffa Credo), kept in an inverted nycthemeral cycle for diurnal schizogony.

**Monoclonal antibodies.** Hybridomas secreting PMMSA specific monoclonal antibodies (mAbs) 1–7, 9–10, 50 and 52 were generated in this laboratory. Spleen cells from hyperimmune BALB/c mice were fused with myeloma cell-line NSO/U [37]. Hybridoma cultures producing antiplasmodial antibodies were identified by indirect immunofluorescence (IIF) and cloned by limiting dilution. Ascites fluid from pristane (2,6,10,14 tetra-

methylpentadecane; Aldrich)-primed mice was used as the source of mAbs. PMMSA-specific mAbs were identified on the basis of their surface reactivity with purified merozoites in suspension (IIF) and immunoprecipitation of an approximately 250-kDa antigen. Mice were made hyperimmune by nivaquine treatment at a parasitemia of 5–25% followed by two more parasite challenges ( $10^7$  infected red cells per mouse) at three-week intervals.

PMMSA-specific mAbs 12.3, 12.11, 12.12, 12.15, 12.17, raised against a cloned *P. c. chabaudi* isolate (isolate CB) [38] were kindly provided by D. Walliker (University of Edinburgh, U.K.). PMMSA-specific mAbs H98 and H100 were a generous gift from M. Hommel (University of Liverpool, U.K.) and had been raised against clone PC-7 of the *P. c. chabaudi* isolate IP-PC1 [39].

**Preparation of cDNA and genomic libraries.** Parasitized blood was collected when infection reached 30–50% and parasites were predominantly at schizont stage. Leukocytes were removed from infected blood as described elsewhere [40]. RNA was extracted by homogenization of saponin-liberated schizonts in 6 M guanidinium-HCl/0.1 M Na-acetate, pH 5.2 and centrifugation through a 4.8 M CsCl/10 mM EDTA (pH 8.0) cushion at 35 000 rev./min in a Beckman SW 41 rotor for 16 h. Poly (A)<sup>+</sup> RNA was selected by oligo(dT) cellulose chromatography and cDNA prepared according to the Amersham cDNA synthesis kit protocol. The cDNA was subsequently methylated with *Eco*RI methylase and ligated to phosphorylated *Eco*RI linkers with T4 DNA ligase. This mixture was then cleaved with *Eco*RI and fractionated on a Bio-Gel A-50m column (Biorad). Fractions containing cDNA molecules > 500 bp were ligated to dephosphorylated  $\lambda$ gt11 *Eco*RI arms and packaged in vitro (Packagene, Promega).

For the construction of the genomic library, *Eco*RI/*Xba*I cleaved genomic DNA was ligated into dephosphorylated lambda GEM-2 *Eco*RI/*Xba*I arms (Promega) and packaged in vitro.

**Screening of the libraries.** Screening of recombi-

nant phag  
cording to  
screening  
the 'Multi  
sham) and  
N (Amer  
[42]. Fina  
ditions (10  
'Na-citrate

**Sequencing.**  
cloned into  
tors. Plasm  
erated tha  
deletions (c  
III digesti  
ing was (c  
or double  
dideoxy ch  
[43] using  
nase, USB  
in the cod  
tially in t  
assisted st  
facilitated  
(Intelligen

**Epitope mapping.**  
cDNA  $\lambda$ gt  
pressing p  
toothpick  
imately 10  
top agaros  
*Escherichia*  
diameter L  
Plates were  
sequently  
ter disk, s  
propyl  $\beta$ -D  
ter and in  
37°C. Nitr  
urated over  
HCl pH 8.0  
taining 1%  
TBST con  
cites fluid  
several tim  
1 h with T  
line phospt

med mice was PMMSA-specific. Analysis of their surzoites in suspension of an approximate made hyper-parasitemia of parasite challenges at three-week

12.11. 12.12. cloned *P. c.* 8] were kindly y of Edinburgh. 198 and H100 mel (University raised against isolate IP-PC1

omic libraries. when infection were predominate. ytes were re-ibed elsewhere. ogenization of guanidinium-centrifugation DTA (pH 8.0) ckman SW 41 is selected by y and cDNA m cDNA syn-subsequently and ligated to T4 DNA lig-1 with *Eco*RI .50m column IA molecules rylated  $\lambda$ gt11 (Packagene.

genomic li-c DNA was bda GEM-2 packaged in

g of recombi-

nant phages with monoclonal sera was done according to Huynh et al. [41]. For hybridization screenings DNA probes were radiolabeled using the 'Multiprime DNA Labelling System' (Amersham) and hybridized to plaque blots on Hybond-N (Amersham) according to standard protocols [42]. Final washes occurred under stringent conditions (10 min at 65°C in 15 mM NaCl/1.5 mM Na-citrate).

**Sequencing.** Inserts of selected clones were subcloned into M13mp18 or pUC 18 sequencing vectors. Plasmid or M13 subclones were then generated that contained progressive unidirectional deletions of each insert by controlled exonuclease III digestion (Erase-a-Base, Promega). Sequencing was done on either single-stranded (M13) or double-stranded (pUC18) templates by the dideoxy chain-termination method of Sanger et al. [43] using modified T7 DNA polymerase (Sequenase, USB). Both strands were entirely sequenced in the coding and 3' untranslated areas and partially in the 5' untranslated region. Computer-assisted storage and analysis of sequence data was facilitated using the PC/GENE software package (Intelligenetics).

**Epitope mapping.** Plaque-purified recombinant cDNA  $\lambda$ gt11 phages 72, 100, 46 and 452, expressing parts of *P. c. chabaudi* PMMSA were toothpicked from suspensions containing approximately  $10^7$  plaque forming units  $\text{ml}^{-1}$  onto the top agarose layer (containing approximately  $10^9$  *Escherichia coli* Y1090 [41] bacteria) of 90 mm diameter LB agar plates ( $100 \mu\text{g}$  ampicillin  $\text{ml}^{-1}$ ). Plates were then incubated at 42°C for 3.5 h, subsequently overlaid with a dry nitrocellulose filter disk, saturated previously with 10 mM isopropyl  $\beta$ -D-thiogalactopyranoside (IPTG) in water, and incubated for an additional 3.5 h at 37°C. Nitrocellulose membranes were then saturated overnight with TBST buffer (10 mM Tris-HCl pH 8.0/150 mM NaCl, 0.05% Tween-20) containing 1% BSA and then incubated for 1 h with TBST containing mouse serum antibody or ascites fluid at a 1/200 dilution. Filters are washed several times with TBST and then incubated for 1 h with TBST containing affinity-purified alkaline phosphatase conjugated goat anti mouse IgG.

antibodies (dilution 1/1000). After a final round of washes substrate (0.05  $\text{mg ml}^{-1}$  5-bromo-6-chloro-3-indolyl acetate, 0.1  $\text{mg ml}^{-1}$  nitro blue tetrazolium in 10 mM  $\text{MgCl}_2$ /100 mM NaCl/100 mM Tris-HCl, pH 9.5) is added to the filters and the enzymatic reaction is stopped after 5–15 min by rinsing the membranes with water. The relative position of cDNA expression clones 72, 100 and 46 in the open reading frame was determined by sequence analysis of the ends of their inserts. Clone 452 was positioned by restriction mapping.

## Results

**Cloning strategy.** cDNA expression libraries ( $\lambda$ gt11) were screened with a mixture of PMMSA-specific mAbs. Several positive clones were detected among which clones 100 and 46 were selected for sequence analysis. Clone 46 was later shown to contain a cloning artefact. This involved the fortuitous ligation of a PMMSA-specific cDNA molecule to an unrelated cDNA molecule (dotted line in Fig. 1). This conclusion was based on Southern blot analysis and comparison with the *P. y. yoelii* YM PMMSA sequence and was confirmed by sequencing genomic clone RX4. Sequence data from the nonspecific part of clone 46 were discarded for this study. Clone R16 was obtained by rescreening cDNA libraries with a radiolabeled DNA fragment that originated from the 3' end of the clone 100 insert.

Clone 100 insert hybridized on Southern blots of *Eco*RI cleaved genomic DNA to a single band of approximately 12 kb, whereas a single 5.4-kb band was detected on Southern blots




Fig. 1. Cloning strategy for the p199 gene. Clones 46, 100 and R16 are  $\lambda$ gt11 cDNA-expression clones. Clone RX4 is a genomic  $\lambda$  clone. The section of clone 46 that originated from a fortuitous ligation event is represented by a dotted line.

of *EcoRI/XbaI* cleaved genomic DNA (data not shown). This 5.4-kb band was cloned by screening a genomic library of *EcoRI/XbaI* cut genomic DNA ligated into  $\lambda$  GEM2 vector with radiolabelled clone 100 insert. In this way genomic clone RX4 (insert size = 5.4 kb) was obtained.

**Sequence analysis.** The inserts of overlapping clones RX4, 46, 100 and R16 were subcloned into M13mp18 or pUC18 vectors and partially or totally sequenced. Together these 4 inserts spanned a region of 6409 bp. This sequence is presented in Fig. 2. One major open reading frame can be found, starting at nucleotide 667 and terminating with a stop codon at position 6022. This codes for a protein of 1785 amino acids (AA) with a calculated  $M_r$  of 198 886. The A+T content of the sequence is high with an average of 67% in the coding region and 83% in the 5' and 3' untranslated region. The A/T ratio of the coding strand is 1.74. This biased A/T ratio of mRNA sense strands appears to be a general phenomenon for malarial genes [44].

The encoded protein has many of the basic features of other PMMSA. At the aminoterminal end a putative signal peptide (residues 1–19) is present while a stretch of hydrophobic amino acids, probably functioning as a membrane anchor sequence, is found at the C-terminus (residues 1765–1785). Ten out of a total of 20 Cys residues are located in the last 110 AA of the protein. Eight potential *N*-glycosylation sites are present. Several tandem repeat oligopeptides, mostly incompletely conserved, are scattered throughout the protein. Analysis at the nucleotide level shows that the individual repeat units are clearly related. Most conspicuous is a stretch of incompletely conserved 7\*6 AA starting at residue 324. Although the other tandem repeat structures are not so extensive, they do occur in many different types. In addition to the above mentioned hexapeptides, tri-, tetra-, penta- and heptapeptide tandem repeats can be observed. Also, a stretch of 7 consecutive alanine residues and a string of 7 consecutive aspartic acid residues are present in the sequence. These can be considered as monocodon repeats.

**Comparison to PMMSA of other species.** The *P. c. chabaudi* PMMSA sequence (p199) was aligned

to known PMMSA sequences of other species using the PALIGN program (PC/GENE). With *P. y. yoelii* YM PMMSA (p197) an overall homology of 69% was detected at the AA level. This homology is not equally distributed along the protein sequence but is clustered in large zones of high homology interspersed with 4 areas of very poor homology (Fig. 3). Large insertions and/or deletions have occurred in these areas. Interestingly, all repetitive sequences are found in these regions. The 20 Cys residues present in both proteins are completely conserved. Alignment with either *P. falciparum* PMMSA allelic sequence (isolates K1 and MAD20) displayed 33% overall homology. This homology is almost exclusively confined to the conserved and semiconserved blocks, as defined by Tanabe [16] (data not shown). The few patches of high homology that occur in the variable blocks are very often also conserved between the 2 *P. falciparum* alleles. The degree of homology appears to be as high with the semi-conserved as with the conserved blocks. From this alignment it is also clear that the 4 divergent areas in the p199/P197 comparison 'coincide' with the *P. falciparum* variable blocks 4, 8, 10 and 14 (as defined by Tanabe).

A crude epitope map of p199 was established by screening the reactivity of a battery of 18 *P. c. chabaudi* PMMSA specific monoclonal antibodies with a set of overlapping cDNA expression clones in  $\lambda$ gt11. Fig. 4 summarizes the results. The major conclusion from this analysis is that all monoclonals seem to map to the central third part of the molecule and that none is binding to the hexapeptide tandem repeats. This area does not seem to be very immunogenic. These data also indicate that carbohydrate moieties do not play a major role in the immunogenicity of the molecule.

## Discussion

In this study we present the complete primary structure of *P. c. chabaudi* PMMSA (p199) based on the DNA sequence of the corresponding gene. P199 exhibits similar characteristics to other PMMSA. It has a calculated  $M_r$  of 199 000, shows putative signal and membrane anchor sequences and a clustering of Cys residues in the last 120 AA.

Although the predicted molecular weights for

species us-  
). With *P. y.*  
II homology  
This homol-  
e protein se-  
of high ho-  
ry poor ho-  
or deletions  
gly. all rep-  
egions. The  
is are com-  
ier *P. falci-*  
tes K1 and  
ology. This  
ned to the  
as defined  
w patches  
ble blocks  
ie 2 *P. fai-*  
y appears  
s with the  
it is also  
199/P197  
rum vari-  
Tanabe).  
established  
f 18 *P. c*  
ntibodies  
on clones  
The ma-  
II mono-  
urt of the  
lexapep-  
m to be  
ate that  
r role in

te pri-  
(p199)  
spond-  
tics to  
99000.  
or se-  
in the  
its for

```
10 20 30 40 50 60
TCTAGATAATATATTTTGTATGCAATGCTAAATTAATATACATATATTAATAGATT
70 80 90 100 110 120
TGTCCGAATCTTTATGTGTGCAATTAATTTTAAATAAATATATCCATATACCAT
130 140 150 160 170 180
TATTAATTTGTGTATACCGTAAATATTTTATTTTAAAGCAATTTTCCTTAAATATAT
190 200 210 220 230 240
TTTTTGTATCATTTTATTTTGTGAAATTCGAGCATATAAATATATATTAC
250 260 270 280 290 300
ACTTTATAAATTTTATACACATTTGTATTTTATTTATATATATTTTAAACAT
310 320 330 340 350 360
TTTTATTTGTAAATGATATGATCAATTAATAAATAAATAACATATAATAGTATAAT
370 380 390 400 410 420
TTTTTTGTAGATATATAAATATATGATTTTATTTTATAGTAAGTTAAAGTGT
430 440 450 460 470 480
ATTATATATGATTTTGTTTTAAACAGAACGAAATAGAAATAACACATATAAATCTAT
490 500 510 520 530 540
ATATATATGTGTAAATAGTGTATGTATATATTTTCAACATTAATGATATAATGTG
550 560 570 580 590 600
AACTTCAATATTTTATACCAATAGTACTAATATAAATGCAAAAGTAATGTACC
610 620 630 640 650 660
TTTCTGTGTATTAATTTTACGATTAATTTATTTTCCACTGTGTATATAGTTAAGTTCC
670 680 690 700 710 720
TTGAAATAGAGCGATTCGAGCTTTTGTCTTTTCTTTTCTTATATATATGCAAA
730 740 750 760 770 780
TCTGAAACATTAAGCAATTTTCAATGATCTGTTTCAATAGTTAGAAAGTTAGAAATTA
790 800 810 820 830 840
TCAATGAGAGGATTAAGCAATTTTCAAAAGTCAAGTAATTTGTAATGCAATCAACA
850 860 870 880 890 900
SVEGLELFGKRSQVIVNAOSP
910 920 930 940 950 960
GAAACACCTTTGTATGCAATTTACAAACCTGAAATTTGCAAAAGTACCAACATTTAT
970 980 990 1000 1010 1020
ETTFVDFFTNPFPAOKLOPFI
1030 1040 1050 1060 1070 1080
TTAAATTTGAGAAATAGGATTTACAGAAACAGAGTAGTCATTTAATAAAGCT
1090 1100 1110 1120 1130 1140
LKFEEELGTYEOTZELVNLKT
1150 1160 1170 1180 1190 1200
TTAGGCCAGCAATTAATTTTACTATGAGCTGCTAGAGATAAATGAAATGATGTCT
1210 1220 1230 1240 1250 1260
LMHAINEFTYDVLDRDLNDNC
1270 1280 1290 1300 1310 1320
GCAAAATACATTTTGTAAATTCCTGAACTCTTAAATTTAATGTGAGCAATCGAAATG
1230 1240 1250 1260 1270 1280
AHNECEIPEERLKNVSEKH
1290 1300 1310 1320 1330 1340
CTTAGGAAGTTGTCTTGGCTATAGAAAGCAATTTGAAATATTCAGATGATCTGTA
1350 1360 1370 1380 1390 1400
LKRVLVQYRKPIENIODDLV
1410 1420 1430 1440 1450 1460
AAATAGAGAAATATATGCAAAATTAAGCACTGCTGAAACCTTAAACCTTTAT
1470 1480 1490 1500 1510 1520
KLESTIARNKATAETLTLI
1530 1540 1550 1560 1570 1580
ACTGAAAGCAAAATAACACTGAAAGCAAGATGCAAGATATAATTTGCT
1590 1600 1610 1620 1630 1640
TERKKITPSEETDCHDTC
```

```
1330 1340 1350 1360 1370 1380
GACAACTAAATATGAAAGAAAAGCAATATATCAAGCTATGACAAATGTTATTT
1390 1400 1410 1420 1430 1440
DNTETGRRKAIYQAHYRVIF
1450 1460 1470 1480 1490 1500
TACAAAAGCAATTAAGTAAATAAATAAATCATGAAAGTCTTGAAGAAAGAGTTGCT
1510 1520 1530 1540 1550 1560
YKKOLAEIKRKVIEVLEKRV
1570 1580 1590 1600 1610 1620
ACATTAAAGAGCAAGCAATTAAGCAATTTTACAAAGCAATGAAAGTATCAGAGCT
1630 1640 1650 1660 1670 1680
TLERKHEAIKPLLOQIEAIRG
1690 1700 1710 1720 1730 1740
CCACTGCTGCTCACTGAAAGCAATTAAGTACAGAAAGCAAGCAAGCAAGCAAGCA
1750 1760 1770 1780 1790 1800
PPAVTEGQIATZGSSSEETKQ
1810 1820 1830 1840 1850 1860
1570 1580 1590 1600 1610 1620
AATAGTACAGAAATCATCTAACCAAAAGCACTACTACTACAAAGCTTTACAAACCA
1630 1640 1650 1660 1670 1680
HSESSMTKTTTDTKAVTTQ
1690 1700 1710 1720 1730 1740
ACCOCTACTAAGCACTGCTACAGAAACAAATATGCTACAGAAACAAATGCTGCA
1750 1760 1770 1780 1790 1800
TATKATGCTETNTCTETNTCT
1810 1820 1830 1840 1850 1860
GAAACAAATCTGCTGCAAGCACTACTGCTGCAAGCACTACTGCTGCAAGCACT
1870 1880 1890 1900 1910 1920
ETNTATGTTTATGTTTATGTTTATGTTTATGTTTATGTTTATGTTTATGTTTAT
1930 1940 1950 1960 1970 1980
CCTACTGCTGCTGCAAGCACTACTGCTGCAAGCACTACTGCTGCAAGCACTACT
1990 2000 2010 2020 2030 2040
PTVTEPVQVPAVQVLTSEER
2050 2060 2070 2080 2090 2100
OCAAAAAATAGCTGAACTTTATGCTCAATTAAGCAAAATGCAAAATGCAAAATG
2110 2120 2130 2140 2150 2160
ARKIAELYAQIKKIAKTIK
2170 2180 2190 2200 2210 2220
AATTAGAGCAATTTGCTGCAATTAAGCAAAATGCAAAATGCAAAATGCAAAATG
2230 2240 2250 2260 2270 2280
NLDGQIFVDFVLELYRK
2290 2300 2310 2320 2330 2340
ARTGAAAGCTGCTGCAAGCACTACTGCTGCAAGCACTACTGCTGCAAGCACTACT
2350 2360 2370 2380 2390 2400
KESCHSTSSCAKNTPTV
2410 2420 2430 2440 2450 2460
CCATTAAATGCTGCAATTTATGCTGCAATTTATGCTGCAATTTATGCTGCAATTT
2470 2480 2490 2500 2510 2520
PLNVRYFPBGISYPLTEEVV
2530 2540 2550 2560 2570 2580
2050 2060 2070 2080 2090 2100
AGCAAAATGCTGCAATTTATGCTGCAATTTATGCTGCAATTTATGCTGCAATTT
2110 2120 2130 2140 2150 2160
SKIAHNAATTTTGGDLTNVDN
2170 2180 2190 2200 2210 2220
ACAGCAATGCAAGCACTACTGCTGCAAGCACTACTGCTGCAAGCACTACTGCT
2230 2240 2250 2260 2270 2280
TATZEDLTTHNQARKNLKA
2290 2300 2310 2320 2330 2340
ATTAAGCAAAATGCAAGCACTACTGCTGCAAGCACTACTGCTGCAAGCACTACT
2350 2360 2370 2380 2390 2400
TKRKIEAESQKLVGLKDDTD
2410 2420 2430 2440 2450 2460
ACTAACTGCAAGCACTACTGCTGCAAGCACTACTGCTGCAAGCACTACTGCT
2470 2480 2490 2500 2510 2520
TKLAAPNQKTPFEREAKRF
2530 2540 2550 2560 2570 2580
2290 2300 2310 2320 2330 2340
TATGAACTCAATTTAGAAATTAATTTACTGCTGCAATTTTGGCAATTTTAAACAAA
2350 2360 2370 2380 2390 2400
YESKERNKLTDTDFDKTR
2410 2420 2430 2440 2450 2460
AGAACTGAAATATGCAAGCACTACTGCTGCAAGCACTACTGCTGCAAGCACTACT
2470 2480 2490 2500 2510 2520
ATEIHNKAAALVGCETONTQ
2530 2540 2550 2560 2570 2580
CAACTCAATTAATTAATTAATTAATTAATTAATTAATTAATTAATTAATTAAT
2590 2600 2610 2620 2630 2640
QLHNKLNKQDNYLQDYGDKR
2650 2660 2670 2680 2690 2700
GAAATGTTTAACTGAAATTTGAAATTTTAAACAAAATGCAAAATGCAAAATG
2710 2720 2730 2740 2750 2760
EIVNTEIEYFSNKRSLQYH
```

2530	2540	2550	2560	2570	2580
ATTAAATAGATTAAGCAATCTGTTTCAAGCAAAACAAATATATATAGTTCGATCAAAACAT					
I M R L A H A V G A R O N I L V A S R H					
2590	2600	2610	2620	2630	2640
ATTCACCTTTCAACACTCTTTTGAATTAACAATCCAAAAATGTTTATTAACAACAACTATTT					
I P L S T L V E L O I Q R B L L T K L I					
2650	2660	2670	2680	2690	2700
GACCAATTAATAAAGCTGATTTTCTTAATAAAGCTCACTTAATAAGCAAGATATAC					
G O L M K T E F E L R N A W L K D E I Y					
2710	2720	2730	2740	2750	2760
GTTCACCAAACTATGCTAAGAAGGAAACAGAACCACTACTACTTAATAGCTATAAAT					
V P O T Y T O R E K C P F Y L I A I R					
2770	2780	2790	2800	2810	2820
AAAGAAATATGACAGACTTGCAGAAATTTATTTCTTAAATATGATGATATGTAAGAAGAG					
X E I D R L A K F I P K I D D H I E R E					
2830	2840	2850	2860	2870	2880
AAACAAAAATGGAAACAGACATGTAGCTACCGGAGAACTGTACAGACGCTCTTCTGCG					
X Q R H E Q E H V A T O E S E C A S S A					
2890	2900	2910	2920	2930	2940
TCTGTACTGGATCATCTCCACAAAGCCACATCAACAAAGCACCACCGCTTCCAGCTGCA					
G O T G S B T E T T S Q T A P A V P A A					
2950	2960	2970	2980	2990	3000
CCCGCACCAACAGAAAGCGAAAGAAAGGACAGAAATCAACAGAGAAAGACCCAGACACA					
P A P A C K A R E G T S B T E E T P A A					
3010	3020	3030	3040	3050	3060
TCAAAACACGACGAAGTGTGACCATCCACAGGTGCAACACCCCAACAGAAACAGAGACT					
S R P A E G A A S T G A T T P T E O C A					
3070	3080	3090	3100	3110	3120
GCACCAACAGAACAGAGACCAACCTGCGACACTTAAACACCAAGCAGAGGTACCAACA					
A P T E O E A O P A A P E T P A S V P A					
3130	3140	3150	3160	3170	3180
CCAACACGCGCTGCGCTCCAGACCACTCCAGCGCCAGCCAGCAGCAGCCGCAAAACCACTT					
P T T P A A P A T P A A P A P A R P V					
3190	3200	3210	3220	3230	3240
ATGACAAATATATATACCTTGAATAATTAAGAATTTTTCAGATCTCATATGCAATGCT					
M T R L Y T L E K L K R F L A P S Y A C					
3250	3260	3270	3280	3290	3300
CATAAATATGTTTATTATCAAAAGCTTACCATTAAACAAAGATGCTTTAAGCAAAATATGCT					
R K Y L L Q N S T I P R D A L S K Y A					
3310	3320	3330	3340	3350	3360
CTTACACACAGAAAGATATAAATAGAACATATTAAGAGATGCTGAATAGATGTATTA					
L T P E E D K I R T L R C E L D V L					
3370	3380	3390	3400	3410	3420
TTAGCTATTCAAAATATGCTTACTATGTATTCTATTATGAAGATGTAGTTGTATGTT					
L A I O N H M P T H Y S L Y B N V D G					
3430	3440	3450	3460	3470	3480
TTACAAAGAACTTACAGTAAATATGAAGAAAGATGATGTATCATATATATACCTTA					
L O N I Y T B L Y E K E M H H Y T N L					
3490	3500	3510	3520	3530	3540
AAAGATAAAACCCAGCTGTTAAAGCTTTATTAAGTAAAGCTGGCGTATTGATCCAGAA					
K D R N A P V A R L L V K A G V I D P E					
3550	3560	3570	3580	3590	3600
CCAGTAGCGCCAGCAACCAAGCACTACCAAGCAGCAAGCAAGCTGCACAGAAAGCTCAGCAGAA					
P V A P T P A V P A P E T A P E T A P E					
3610	3620	3630	3640	3650	3660
ACTGCACAGAAACCAACAGCAAGAAAGCTCCACCAACCAACAGAAATGGCCACAGCAGCA					
Y A P E T P A Q E A P Q O P E D A Q A P					
3670	3680	3690	3700	3710	3720
GAAGCAGCACTGAACACCAACCAAGCAGCAATCGGATCAACAGCAACCAACCAAGCAAGAA					
S A A T E T T T P A S S A S T E P T R K					

3730	3740	3750	3760	3770	3780
GCACCTACACCAACACCCACATCTGTAAACAGTATACACAGAGGAGAACACACCCACGCA					
A P T A T T P T S E Y T V T Q E G T T P A A					
3790	3800	3810	3820	3830	3840
CCAAAACACACAGAGGAGACATATCATCAGCACCCAGCACACACCCAGCCAGCAAAACCA					
F R R O E G A S S S A P A Q P A P A R P					
3850	3860	3870	3880	3890	3900
GCACCTCCACAAACAGTAAACAGCCACATCAACAAACOTTGTGAAGGAAGTACTCAAGTAAAG					
A P A Q T V T G Q S T N V E G S T Q V R					
3910	3920	3930	3940	3950	3960
GCAGAAAGTGAAGACGAAATGTTATGTCGATGATTTCGAATAGACATTTTTCACAAATCT					
A E S E D E H F V D D F E V D N F T Y K E					
3970	3980	3990	4000	4010	4020
TACTTACACCAAGTGTATGGAATAATATGCAATTCATGATTTTATATAAATCTAAAAAA					
T L O Q V D Q H N T T Q T D F I K E K K					
4030	4040	4050	4060	4070	4080
GAATTATCTAATCATTTGACCCCTGAAAAAAGTTAAACCAATATATCTTGTATGTGACAC					
E L I N A L T P E K V N O L Y L D I A H					
4090	4100	4110	4120	4130	4140
TTAAAGGATATATGACACCAATCTACTATAATCTGTTATATAAATAAATTTAAATAGAA					
L F E L S E N T Y T T T T T T E K S L K L E					
4150	4160	4170	4180	4190	4200
AGATTTATCTAAAAACATGAACCAATTTGAAGCAGCTATACCAAAAAGTTTAAAGAAATTAAC					
R L Y G R H E Q I E A A N O K V K E I S					
4210	4220	4230	4240	4250	4260
GTATTAAATCCGATTTATAAAAAGAAAAATATATTTAAATGTCATTTTATGTATTA					
V L K E R L L K R K K Y I N G T T F Y V L					
4270	4280	4290	4300	4310	4320
TCTGGTTTCCAAATTTCTTTAAACAGAGAGAGAGAGCTGAAAAGCTATATGTATATAC					
S G F A N F F N K R R R B A E K Q Y V D H					
4330	4340	4350	4360	4370	4380
GCATTAATAAAATACTGATATGTTATTGAAATCTCAAAAGCTCTGATGAAATTTTACT					
A Z K R T D M L L K T Y K A R S K T F T					
4390	4400	4410	4420	4430	4440
TCTGAAGCTTCTCTTTAAAAACATTAACATAACATCAATTCACAGAGAGACCCATCTAC					
S E A V P L F T L T L R T S I D R E A N Y					
4450	4460	4470	4480	4490	4500
TTTAAATTTGAAAATTTCAAGACATACACGGGATAGAAATTAAGATTTAAAGAAAAATTT					
L K I E K F R A Y S R L E L L L K K K N I					
4510	4520	4530	4540	4550	4560
AACCTAGGAAAAAGAAATATACATATGTTATCTGTTGTTTACACATGTTATTTGAAGAA					
N L G K E R I Y T V S G O L N H V F E E					
4570	4580	4590	4600	4610	4620
TTTAAAGAAATTTTAAAAATAAAGGTTATACCGGAAAAATTAACCTTGAANAATGCTCTCT					
F K E L L K N K G Y T G E T H P E N A P					
4630	4640	4650	4660	4670	4680
GAAGTTATCAAGGCATTCGACCAATATAAAGAAITACTTCCAAAGGAGCAACCAATCCA					
Z V I R A F E Q E K S L L P K G A T T P					
4690	4700	4710	4720	4730	4740
GCTCAAGTTATTGACCTCTGATGTTCTGTCGAGCCGACACTACAGACGCCGACAGCTAC					
A P V V A P V V A P A P A T A A P A A D					
4750	4760	4770	4780	4790	4800
GCACAGTACACGACAGCCGACGCGGACGCGGACCTCAGATCAGGATCAGCAGCCACACCA					
A P V P A A A A A A A S G S G A T T T					
4810	4820	4830	4840	4850	4860
GAAGGAGAGCCGCTACTACATAGTATTCAGACGACGCGGATATGATGATGATGATGATGAT					
E G A A T T V V A S S D H D D D D D					
4870	4880	4890	4900	4910	4920
GATATGATCAAAATTCGAAATGCTCAATCCACAGAGAGAGGATTAAGAAATATCTTGAT					
D N O G I A N A Q S T D S E V R D I L D					

Fig. 2. Nuclear reading frame are doubly u

PMMSA of *yoelii* (197 kDa) [14] molecules migrate more slowly than those of *falciparum* between apparent  $M_r$  has been determined by immunoprecipitation and antigenic analysis. It has been announced by the authors that the presence of the proper antigens are related to p199 and p190 of *P. falciparum*.




Fig. 2. Nucleotide sequence of the p199 gene of *P. c. chabaudi* and the deduced amino acid sequence of the major open reading frame. Putative signal and membrane anchor sequences are underlined by a dotted line, whilst tandem repeat structures are doubly underlined. Cys residues and potential N-glycosylation sites are denoted by asterisks and '+' signs respectively. Nucleotide 186 is as yet undetermined.

PMMSA from *P. c. chabaudi* (199 kDa), *P. y. yoelii* (197 kDa) [24] and *P. falciparum* (188–196 kDa) [14–16, 18–19] are rather similar, PMMSA molecules from both rodent malaria parasites migrate more slowly on SDS-PAGE than their *P. falciparum* counterparts [45]. A discrepancy between apparent  $M_r$  on SDS-PAGE and predicted  $M_r$  has been generally observed for malarial repetitive antigens [46]. It is thought that the pronounced hydrophilic character of these antigens and the presence of repeat structures interfere with the proper binding of SDS so that these antigens are retarded in SDS-PAGE. However both p199 and p197 are neither more hydrophilic than *P. falciparum* PMMSA nor do they contain dra-

matically more repeats (p197 actually has less). The most plausible explanation would therefore be that PMMSA of rodent malaria parasites are more extensively glycosylated than *P. falciparum* PMMSA. This glycosylation could occur at one or more of the 4 potential N-glycosylation sites that are conserved between *P. c. chabaudi* and *P. y. yoelii* (residues 662, 866, 1106 and 1640 in p199). On the other hand Holder and Freeman were unable to find experimental evidence for glycosylation in *P. y. yoelii* PMMSA [9]. The physicochemical basis of this retarded migration in SDS-PAGE remains therefore unclear.

Southern blots probed with clone 100 insert only showed 1 band. Also not a single nucleotide

238

P199	- MKAIGLLFSFVFFAIYCKSETIGVYNLDLVHKLKLEELSVEGLELFQKSQVIINA	-55
	: : : : : * : : : : : : : : : : : : : : : :	
P197	- MKVIGLLFSFVFFAICKCKSETIEVYNDLIQKLEKLESLSVDGLELFQKSQVIINA	-55
P199	- QSPETVPDPFTNPEFAQKLQPFILKFEEFGTEQTELVNLIKTLGPNKYGLKYLI	-110
	: :	
P197	- TQPTETIDPFTNNHFAQQVQDFVTKEFEGLGTEQTELVNLIKALTPNRXYGVKKYLI	-110
P199	- ESKEEFNELMHAINFYDVLDRDKLNDMCANNYCEIPEHLKINVEEIEMLKKVVVG	-165
	: : : : : : : : : : : * : : : : * : : : : : : : : : :	
P197	- ESKEEFENGLMHAINFYDVLDRDKLNDMCANNYCEIPEHLKISEEETEMLKVVVG	-165
P199	- YRKPIENIQDDLVLKLEEIARNKATAETLNTLITEETKKITPEEETDCNDTNCND	-220
	: : : : : : : : : : : : : : : : * : : : *	
P197	- YRKPIENIQDDIEKLEIYIERNKETVAALNALIAEETKKIQPEGNECDNASCD	-220
P199	- TKYGKKKAIYQAMYNVIFYKKQLAEIKKVIEVLEKR VATLKKN EAIKPLLQOIFA	-275
	: :	
P197	- DKYNKKKIYQAMYNVIFYKKQLAEIQKVVLEVLEKR VSTLRKKNDAIKPLWQQIEV	-275
P199	- IRGPPAVTEG-QIATEGSSEETKQNSTESSNTKTITTDKAVTTQTATKATGTETN	-329
	: : : : : : : : : : : : : : : :	
P197	- LNAAPVTAETQIVTGG-----QSSTEP-----GSGGS	-303
P199	- TGTEINTGTETNTATGTTTATGTTTATGTPV-----TEPVQVPAVQVLTE	-375
	: : : : : : : : : : : : : : : :	
P197	- SASGTSSSGQASAGTGVEQANTVASVTVPISVGNGEASTNPQTAQVQPVTILT	-358
P199	- EEKAKKIAELYAQIKEIAKTIKFNLDGIFVDPVELEYKKEKKNESCH-STSSCH	-429
	: : : : : : : : : : : : : : : * : : : *	
P197	- EEKQKKIAGLYAQIKEIAKTIKFNLEGIFVDP IELEYFKKEKKESCNLSTSSCK	-413
P199	- KNKTPETVIPLNVRYPNGISYPLTEEVSYSKIAHNAAETTYGDLTNVDNTAIED	-484
	: :	
P197	- KKNASETIIPLTIRYPNGISYPLPENDVYNKIANNAAETTYGDLTHPDNTPLTGD	-468
P199	- LTNEQARKNLIAIKKKIEAEQKLVELKDDYDTKLAAFNGQKTPFKEAAKFFY	-539
	: :	
P197	- LATNEQARKDLIAIKKKIKAEKKLETILKTNYDNKLTENQKTPFKEAAKEFY	-523
P199	- ESKFRNKLTTFDDFKTKRTEYMNKKAALVGCEYGNTQQLINKLNKQNLNYLDY	-594
	: : : : : : : : : : : : : : : * : : : : : : : : : :	
P197	- ESKFRNKLTSEIFEKFKTRDEYMTKKTELNTCEYGNTKELINKLNKQNLNYLDY	-578
P199	- GLRKEIVNTEIEYFSNKKSELQYNINRLANAVQAKONILVASKHIPLSTLVELQI	-649
	: :	
P197	- SLRKDIISNEIEYFSNKKSELQYNINRLAEAVQAKONVLVASKDVPLSTLVELQI	-633
P199	- QKSLLTKLIEQLNKTEFSLNKAHLKDKIYVPQTYGEGKPEPYLLIAIKKEIDRL	-704
	: : : : + :	
P197	- QKSLLTKQIEQLNKTEVSLNKAQLKDKLYVPKTYGNEGKPEPYLLIAVKKEVDRL	-688
P199	- AKFIPKIDDMIEKEKQKMEQEHVATGESEBQASSAGTSSTETTSQTA-----	-752
	: :	
P197	- AQFIPKIESMIAKEKERMEQGPAITGESEBVPSPSAESSTDRTSOSTSSSSSS	-743



[illegible]

```
P199 - NTKFVVKNPYQLLDNDKKDKQMINLKYAIKGVTEIETATDGIFFNKMIELYK -1537
      : : : : : : : : : : : : : : : : : : : : : : : :
P197 - STNKYVIRNPYQLLDNDKKDKQIVNLKYATRGINEDIETTTDGIKFFNKMVELYN -1523

P199 - PQLNAVNEQIAAIGTEPTDA---EKKKYAIPFEDLKGLYETIILNGAEFSELLQH -1589
      : : : : : : : : : : : : : : : : : : : : : : : :
P197 - TQLAAVKEQIATIEAETNDTNKEEKKKYIPILEDLKGLYETVIGQAEYSEELQN -1578

P199 - KLENYKIEKAGFDILMANLETYIRIDEKLEDFVESAEKNKHIAIALNNLNKSGL -1644
      : : : : : : : : : : : : : : : : : : : : : : : :
P197 - RLDNYKNEKAEFEILTNNLEKYIQIDEKLEDFVEHAENNNKHIAIALNNLNKSGL -1633

P199 - VTEGESKKILAKMLNMDAMDLLGIGSNHVCISTS-TPDNAGCFRYDDGTEEWRCL -1698
      : : : : : : : : : : : : * : : : * : : : * : :
P197 - VGEGESKKILAKMLNMDGMDLLGVDPKHVCVDRDIPKNAGCFRDDNGTEEWRCL -1688

P199 - LGFKKDDGNGRCVADDAPVCNNNNNGGCDKNADCREVENTDRDP SKKIVCTCKEPN -1753
      : : : : : * : : * : : * : : : : : * : :
P197 - LGYKKGE-GNTCVENNNPTCDINNGGCDPTASCQNAESTEN--SKKIICTCKEPT -1740

P199 - PNAYYAGVFCSSSGFMGLSILLIITLIVFNLF -1785
      : : : : : * : : : : : : : : : : : :
P197 - PNAYYEGVFCSSSGFMGLSILLIITLIVFNIF -1772
```

Fig. 3. Alignment of p199 with *P. yoelii* PMMSA (p197). Residues that are part of repetitive structures in both p199 and p197 are in bold face. Positions of Cys residues that are conserved are indicated by asterisks whilst conserved potential *N*-glycosylation sites are denoted by '+' signs. The four highly divergent areas have been boxed.

of difference was observed between genomic clone RX4 and cDNA clones 100 and 46 in the areas where both types of clone have been sequenced (totalling about 1500 nucleotides). Taken together these data provide strong evidence that the p199 gene occurs as a single copy in the genome.

The protein sequence of p199 is very homologous to p197 (69%) and 33% homologous to both *P. falciparum* PMMSA allelic sequences. The same level of homology (31%) had previously been observed between these 2 allelic sequences and p197 [24], indicating that, as might have been expected, *P. c. chabaudi* and *P. y. yoelii* are evolutionary equidistant to *P. falciparum*. The major findings of the PMMSA interspecific comparisons, as described here, are schematically represented in Fig. 5 and can be summarized as follows: (i) p199 and p197 are rather homologous but differ extensively in 4 areas that correspond to *P. falciparum* PMMSA variable blocks 4, 8, 10 and 12. On the contrary variable blocks 2, 6 and 16 are well conserved between p199 and p197. (ii) The homology that exists between p199/p197 and *P. falciparum* PMMSA is situated almost exclusively in the conserved and semi-conserved blocks. The

variable blocks are nearly totally divergent. In these variable areas small patches of homology occur that frequently correspond to stretches that are also conserved between the 2 *P. falciparum* alleles.

It is assumed that the *P. falciparum* alleles evolved in 2 biologically isolated populations that later on merged [16]. The rodent malaria sequence data further illustrate the evolutionary behavior of PMMSA: some areas of the protein are well conserved whereas others are very variable. It is interesting that whereas 7 hypervariable areas are apparent when comparing the 2 *P. falciparum* alleles or when comparing *P. falciparum* PMMSA with the rodent malaria PMMSA, *P. c. chabaudi* and *P. y. yoelii* only diverge profoundly in 4 of these 7 variable blocks. This might indicate that variable blocks 4, 8, 10 and 14 evolve even faster than variable blocks 2, 6 and 16.

The question might be asked as to why some regions in the molecule did evolve very rapidly while other parts changed at a slower pace. Clearly there must be structural and/or functional constraints on the more conserved parts of the protein. These data might however also indicate that the

Fig. 4. Crude reading frame mAbs that react with the 3' region of the *gag* gene is indicated by

divergent zo  
which incre  
types of pos  
are the need  
and immun  
considered t  
invasion of

Fig. 5. Scheme with the PMM into conserved where p199 and MAD20/K homology to 1 been kept unc occurred in so

1537

1523

1589

1578

.644

.633

698

688

753

740




Fig. 4. Crude epitope map of p199. The relative positions of  $\lambda$ gt11 expression clones 72, 100, 46 and 452 to the total open reading frame are indicated. Amino acid numbering is indicated below the bar representing the open reading frame. Names of mAbs that react with each of these expression clones are bracketed. The section of clone 46 that represents the cloning artefact is indicated by a dotted line. The 4 divergent regions in p199 are denoted by chequered blocks. The areas to which various epitopes can be mapped are indicated by open boxes.

p199 and p197  
glycosylation

vergent. In  
homology  
retches that  
*falciparum*

um alleles  
lations that  
a sequence  
behavior of  
well con-  
e. It is in-  
areas are  
*iparum* al-  
PMMSA  
*chabaudi*  
ly in 4 of  
licate that  
ven faster

why some  
y rapidly  
e. Clearly  
nal con-  
e protein.  
: that the

divergent zones were subject to positive selection which increased the rate of genetic change. Two types of positive selection that might be envisaged are the need to adapt to an evolving vertebrate host and immunological pressure. Since PMMSA is considered to participate in the recognition and/or invasion of red cells [47], the variable blocks,

which differ considerably between different *Plasmodium* species and between the 2 *P. falciparum* alleles, might constitute the domains that mediate this interaction. It must be assumed then that the 2 *P. falciparum* PMMSA alleles evolved to interact with different structures on the human red cell membrane (as proposed by Tanabe et al. [16]) and




Fig. 5. Schematic comparison of the amino acid sequences of the 2 *P. falciparum* PMMSA alleles (strains MAD20 and K1) with the PMMSA amino acid sequences from *P. c. chabaudi* (p199) and *P. y. yoelii* (p197). *P. falciparum* sequences are divided into conserved blocks (open boxes), semi-conserved blocks (boxes with vertical lines) and variable blocks (black boxes). Areas where p199 and p197 are highly divergent are denoted by chequered boxes. Regions that show homology between p199/p197 and MAD20/K1 are indicated by stippled boxes whilst blocks that are highly conserved between p199 and p197 but show no homology to MAD20/K1 are shaded. For clarity the relative sizes of the different blocks in the *P. falciparum* alleles have been kept unchanged in the p199/p197 block diagram. It should be noted however that due to insertions/deletions that have occurred in some variable blocks of p199/p197 the relative size of these blocks is not accurately represented. The numbering of the different blocks is according to Tanabe et al. [16].

also that *P. c. chabaudi* and *P. y. yoelii* PMMSA recognize different groups on the mouse red cell. It is however possible that the need to interact with different red cell structures in different hosts can be accommodated by minor AA changes in the more conserved areas.

Alternatively the rapid genetic change in the divergent areas might have been generated by immunological pressure. The repetitive motifs which are found in these areas might be the result of special genetic mechanisms that warrant rapid diversification for the evasion of immune responses. The strain-specific protection shown by PMMSA of *P. c. chabaudi* AS and CB [48] and the isolation of anti-PMMSA mAb resistant lines from cloned *P. c. chabaudi* AS [49] is in keeping with this hypothesis. Interesting also in this respect is the observation that *P. c. chabaudi* PMMSA appears to be even more polymorphic than *P. falciparum* PMMSA since every field isolate out of 15 tested belonged to a different PMMSA serotype (McLean, A.P., Ph.D.-Thesis, University of Edinburgh, 1986).

At the same time these variable regions are probably not very immunodominant. This is indicated by our epitope mapping studies. The first variable area (block 4) is not recognized by any of the PMMSA specific monoclonals while the epitopes of 4 mAbs were shown to map outside of the 4 divergent areas. On the basis of its similar structure (alternating tripeptide repeats) the major repeat area in p199 (residues 324-365) appears to be the homologue to the tripeptide repeats seen in PMMSA of most *P. falciparum* isolates. However the *P. falciparum* tripeptide tandem repeats occur in variable block 2 whereas the p199 repeats are found in variable block 4. The tripeptide tandem repeat area in *P. falciparum* PMMSA is widely polymorphic among different isolates. Work is in progress to determine whether a similar polymorphism prevails in different *P. c. chabaudi* strains.

#### Acknowledgements

This work was supported by a Smith Kline RIT/IWONL contract and by EEC contracts TSD-M-152 and TS-2-0148. We wish to thank Drs. M. Hommel and D. Walliker for providing mAbs and Drs. Jef Seurinck and Gaston Mathijssen (Plant

Genetic Systems) for help with sequencing work.

#### References

- 1 Newbold, C.I. (1985) Parasite antigens in protection diagnosis and escape: *Plasmodium*. In: Current Topics in Microbiology and Immunology, pp. 69-104, Springer-Verlag, Berlin, Heidelberg.
- 2 Hall, R., Osland, A., Hyde, J.E., Simmons, D.L., Hope, I.A. and Scaife, J. (1984) Processing, polymorphism, and biological significance of p190, a major surface antigen of the erythrocytic forms of *Plasmodium falciparum*. Mol Biochem. Parasitol. 11, 61-80.
- 3 Lyon, J.A., Geller, R.H., Haynes, J.D. and Weber, J.L. (1986) Epitope map and processing scheme for the 195 000-dalton surface glycoprotein of *Plasmodium falciparum* merozoites deduced from cloned overlapping segments of the gene. Proc. Natl. Acad. Sci. USA. 83, 2989-2993.
- 4 Freeman, R.R. and Holder, A.A. (1983) Surface antigens of malaria merozoites. A high molecular weight precursor is processed to an 83,000-mol.wt. form expressed on the surface of *Plasmodium falciparum* merozoites. J. Exp. Med. 158, 1647-1653.
- 5 David, P.H., Hadley, T.J., Aikawa, M. and Miller, L.H. (1984) Processing of a major parasite surface glycoprotein during the ultimate stages of differentiation in *Plasmodium knowlesi*. Mol. Biochem. Parasitol. 11, 267-282.
- 6 Heidrich, H.-G., Matzner, M., Miettinen-Baumann, A. and Strych, W. (1986) Immunoelectron microscopy shows that the 80 000-dalton antigen of *Plasmodium falciparum* merozoites is localised in the surface coat. Z. Parasitenkd. 72, 681-683.
- 7 Atkinson, C.T., Aikawa, M., Fujino, T., Tani, L.Q., Hu, G.S.N. and Siddiqui, W.A. (1987) Ultrastructural localization of protective and non-protective *Plasmodium falciparum* proteins using serum samples from vaccinated adult monkeys. J. Parasitol. 73, 1235-1240.
- 8 Holder, A.A. (1988) The precursor to major merozoite surface antigens: structure and role in immunity. Prog. Allergy. 41, 72-97.
- 9 Holder, A.A. and Freeman, R.R. (1984) Characterisation of a high molecular weight protective antigen of *Plasmodium yoelii*. Parasitology. 88, 211-219.
- 10 Boyle, D.B., Newbold, C.I., Smith, C.C. and Brown, K.N. (1982) Monoclonal antibodies that protect in vivo against *Plasmodium chabaudi* recognize a 250,000-dalton parasite polypeptide. Infect. Immun. 38, 94-102.
- 11 Newbold, C.I., Boyle, D.B., Smith, C.C. and Brown, K.N. (1982) Identification of a schizont and species specific surface glycoprotein on erythrocytes infected with rodent malarial. Mol. Biochem. Parasitol. 5, 45-54.
- 12 McBride, J.S., Newbold, C.I. and Anand R. (1985) Polymorphism of a high molecular weight schizont antigen of the human malaria parasite *Plasmodium falciparum*. J. Exp. Med. 161, 160-179.
- 13 Newbold, C.I., Schryer, M., Boyle, D.B., McBride, J.S., McLean, A., Wilson, R.J.M. and Brown, K.N. (1984) A possible molecular basis for strain specific immunity to malaria. Mol. Biochem. Parasitol. 11, 337-347.
- 14 Holder, A.A., Lockyer, M.J., Odink, K.G., Sandhu, J.S., Riveros-Moreno, V., Nicholls, S.C., Hillman, Y., Davey, L.S., Tizard, M.L.V., Schwarz, R.T. and Freeman, R.R. (1985) Primary structure of the precursor to the three major

surf  
Natu  
15 Mac  
Cert  
mon  
of P  
netic  
16 Tan  
(198  
the  
195.  
17 Web  
ation  
gen  
Nuc  
18 Pete  
Woo  
(198  
surf  
Para  
19 Char  
S.E.  
gene  
zoite  
isola  
20 Myl  
sequ  
falc  
5401  
21 Cert  
(198  
face  
lack  
22 Pete  
D.J.  
zoite  
Cell  
23 Del  
Uda  
mod  
stagi  
24 Lew  
ing  
mod  
25 Hol  
agai  
anti  
26 Sidd  
Case  
Kan  
tein  
ium  
301  
27 Hall  
I.A.  
gen  
in b  
28 Perr  
J. ar  
mon  
ual  
29 Che  
J., C  
mun  
cipa

- sequencing work.
- ens in protection diag-  
Current Topics in Mi-  
-104. Springer-Verlag.
- Simmons, D.L., Hope,  
g. polymorphism, and  
major surface antigen  
*Plasmodium falciparum*. Mol.
- J.D. and Weber, J.L.  
ing scheme for the  
of *Plasmodium falciparum*  
ned overlapping seg-  
cad. Sci. USA. 83.
- 83) Surface antigens  
cular weight precu-  
r form expressed on  
merozoites. J. Exp.
- M. and Miller, L.H.  
surface glycoprotein  
in *Plasmodium*  
l. 267-282.
- en-Baumann, A. and  
microscopy shows that  
in *Plasmodium falciparum* mero-  
zoites. Z. Parasitenkd. 72.
- T., Tam, L.Q., Hui,  
Ultrastructural local-  
ization of *Plasmodium falciparum*  
in vaccinated Aotus
- o major merozoite  
in immunity. Prog.
- Characterisation of  
gen of *Plasmodium*
- and Brown, K.N.  
ect in vivo against  
000-dalton parasite
- and Brown, K.N.  
id species specific  
fected with rodent
- 5-54.
- id R. (1985) Poly-  
chizont antigen of  
*falciparum*. J. Exp.
- B., McBride, J.S.,  
n. K.N. (1984) A  
cific immunity to  
37-347.
- .G., Sandhu, J.S.,  
Iman, Y., Davey,  
id Freeman, R.R.  
to the three major
- surface antigens of *Plasmodium falciparum* merozoites.  
Nature. 317, 270-273.
- 15 Mackay, M., Goman, M., Bone, N., Hyde, J.E., Scaife, J.,  
Certa, U., Stunnenberg, H. and Bujard, H. (1985) Poly-  
morphism of the precursor for the major surface antigens  
of *Plasmodium falciparum* merozoites: studies at the ge-  
netic level. EMBO J. 4, 13B, 3823-3829.
- 16 Tanabe, K., Mackay, M., Goman, M. and Scaife, J.G.  
(1987) Allelic dimorphism in a surface antigen gene of  
the malaria parasite *Plasmodium falciparum* J. Mol. Biol.  
195, 273-287.
- 17 Weber, J.L., Leininger, W.M. and Lyon, J.A. (1986) Vari-  
ation in the gene encoding a major merozoite surface anti-  
gen of the human malaria parasite *Plasmodium falciparum*.  
Nucleic Acids Res. 14, 3311-3323.
- 18 Peterson, M.G., Coppel, R.L., McIntyre, P., Langford, C.J.,  
Woodrow, G., Brown, G.V., Anders, R.F. and Kemp, D.  
(1988) Variation in the precursor to the major merozoite  
surface antigens of *Plasmodium falciparum*. Mol. Biochem.  
Parasitol. 27, 291-302.
- 19 Chang, S.P., Kramer, K.J., Yamaga, K.M., Kato, A., Case,  
S.E. and Siddiqui, W.A. (1988) *Plasmodium falciparum*:  
gene structure and hydrophathy profile of the major mero-  
zoite surface antigen (gp 195) of the Uganda-Palo Alto  
isolate. Exp. Parasitol. 67, 1-11.
- 20 Myler, P.J. (1989) Nucleotide and deduced amino acid  
sequence of the gp195 (MSA-1) gene from *Plasmodium*  
*falciparum* Palo Alto PLF-3/B11. Nucleic Acids Res. 17,  
5401.
- 21 Certa, U., Rotmann, D., Matile, H. and Reber-Liske, R.  
(1987) A naturally occurring gene encoding the major sur-  
face antigen precursor of p190 of *Plasmodium falciparum*  
lacks tripeptide repeats. EMBO J. 6, 4137-4142.
- 22 Peterson, M.G., Coppel, R.L., Moloney, M.B. and Kemp,  
D.J. (1988) Third form of the precursor to the major mero-  
zoite surface antigens of *Plasmodium falciparum*. Mol.  
Cell. Biol. 8, 2664-2667.
- 23 Del Portillo, H.A., Gysin, J., Mattei, D.M., Khouri, E.,  
Udagama, P.V., Mendis, K.N. and David, P.H. (1988) *Plas-*  
*modium vivax*: cloning and expression of a major blood-  
stage surface antigen. Exp. Parasitol. 67, 346-353.
- 24 Lewis, A.P. (1989) Cloning and analysis of the gene encod-  
ing the 230-kilodalton merozoite surface antigen of *Plas-*  
*modium yoelii*. Mol. Biochem. Parasitol. 36, 271-282.
- 25 Holder, A.A. and Freeman, R.R. (1981) Immunization  
against blood-stage rodent malaria using purified parasite  
antigens. Nature 294, 361-364.
- 26 Siddiqui, W.A., Tam, L.Q., Kramer, K.J., Hui, G.S.N.,  
Case, S.E., Yamaga, K.M., Chang, S.P., Chan, E.B.T. and  
Kan, S.C. (1987) Merozoite surface coat precursor pro-  
tein completely protects Aotus monkeys against *Plasmod-*  
*ium falciparum* malaria. Proc. Natl. Acad. Sci. USA 84,  
3014-3018.
- 27 Hall, R., Hyde, J.E., Goman, M., Simmons, D.L., Hope,  
I.A., Mackay, M. and Scaife, J. (1984) Major surface anti-  
gen gene of a human malaria parasite cloned and expressed  
in bacteria. Nature. 311, 379-382.
- 28 Perrin, L.H., Merkli, B., Loche, M., Chizzolini, C., Smart,  
J. and Richle, R. (1984) Antimalarial immunity in Saimiri  
monkeys. Immunization with surface components of asexual  
bloodstages. J. Exp. Med. 160, 441-451.
- 29 Cheung, A., Leban, J., Shaw, R.A., Merkli, B., Stocker,  
J., Chizzolini, C., Sander, C. and Perrin, L.H. (1986) Im-  
munization with synthetic peptides of a *Plasmodium fal-*  
*ciparum* surface antigen induces antimerozoite antibodies.
- Proc. Natl. Acad. Sci. USA. 83, 8328-8332.
- 30 Patarroyo, M.E., Romero, P., Torres, M.L., Clavijo, P.,  
Moreno, A., Martinez, A., Rodriguez, R., Guzman, F.  
and Cabezas, E. (1987) Induction of protective immunity  
against experimental infection with malaria using synthetic  
peptides. Nature. 328, 629-632.
- 31 Patarroyo, M.E., Amador, R., Clavijo, P., Moreno, A., Guz-  
man, F., Romero, P., Tascon, R., Franco, A., Murillo, L.A.,  
Ponton, G. and Trujillo, G. (1988) A synthetic vaccine pro-  
tects humans against challenge with asexual blood stages  
of *Plasmodium falciparum* malaria. Nature. 332, 158-161.
- 32 Freeman, R.R. and Holder, A.A. (1983) Characteristics of  
the protective response of Balb/c mice immunized with  
a purified *Plasmodium yoelii* schizont antigen. Clin. Exp.  
Immunol. 54, 609-616.
- 33 Playfair, J.H.L., De Souza, J.B., Freeman, R.R. and Holder,  
A.A. (1985) Vaccination with a purified blood-stage malaria  
antigen in mice: correlation of protection with T cell-  
mediated immunity. Clin. Exp. Immunol. 62, 19-23.
- 34 Majarian, W.R., Daly, T.M., Weidanz, W.P. and Long,  
C.A. (1984) Passive immunization against murine malaria  
with an IgG3 monoclonal antibody. J. Immunol. 132,  
3131-3137.
- 35 Burns, J.M., Jr., Daly, T.M., Vaidya, A.B. and Long, C.A.  
(1988) The 3' portion of the gene for a *Plasmodium yoelii*  
merozoite surface antigen encodes the epitope recognized  
by a protective monoclonal antibody. Proc. Natl. Acad. Sci.  
USA. 85, 602-606.
- 36 David, P.H., Hommel, M., Benichou, J.C., Eisen, H. and  
Pereira da Silva, L.H. (1978) Isolation of malaria mero-  
zoites: Release of *Plasmodium chabaudi* merozoites from  
schizonts bound to immobilized concanavalin A. Proc.  
Natl. Acad. Sci. USA 75, 5081-5084.
- 37 Kearney, J.F., Radbruch, A., Liesegang, B. and Rajewsky,  
K. (1979) A new mouse myeloma cell line that has lost  
immunoglobulin expression but permits the construction  
of antibody-secreting hybrid cell lines. J. Immunol. 123,  
1548-1550.
- 38 Killick-Kendrick, R. and Peters, W. (1978) Rodent Malaria,  
Academic Press, London, New York, San Francisco.
- 39 Hommel, M., David, P.H., Guillote, H. and Pereira da  
Silva, L. (1982) Protection against *Plasmodium chabaudi*  
malaria, I. Vaccination of mice with merozoites and  
Freund's adjuvants. Ann. Immunol. (Inst. Pasteur) 133C,  
57-67.
- 40 Howard, R.J., Smith, P.M. and Mitchell, G.F. (1978) Re-  
moval of leucocytes from red cells in *Plasmodium berghei*-  
infected mouse blood and purification of schizont-infected  
cells. Ann. Trop. Med. Parasitol. 72, 573-575.
- 41 Huynh, T.V., Young, R.A. and Davis, R.W. (1986) Con-  
structing and screening cDNA libraries in lambda gt10 and  
lambda gt11. In: DNA Cloning Vol. I (Glover, D.M., ed.),  
pp. 49-78, IRL Press, Oxford/Washington DC.
- 42 Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982) Molec-  
ular Cloning. A Laboratory Manual. Cold Spring Harbor  
Laboratory, Cold Spring Harbor, NY.
- 43 Sanger, F., Nicklen, S. and Coulson, A.R. (1977) DNA  
sequencing with chain-terminating inhibitors. Proc. Natl.  
Acad. Sci. USA. 74, 5463-5467.
- 44 Weber, J.L. (1988) Molecular biology of malaria parasites.  
Exp. Parasitol. 66, 143-170.
- 45 Holder, A.A., Freeman, R.R. and Newbold, C.I. (1983)  
Serological cross reaction between high molecular weight  
proteins synthesised in blood schizonts of *Plasmodium*  
*yoelii*, *Plasmodium chabaudi* and *Plasmodium falciparum*.

- Mol. Biochem. Parasitol. 9, 191-196.
- 46 Anders, R.F., Coppel, R.L., Brown, G.V. and Kemp, D.J. (1988) Antigens with repeated amino acid sequences from the asexual blood stages of *Plasmodium falciparum*. *Prog. Allergy*. 41, 148-172.
- 47 Hadley, T.J., Klotz, F.W. and Miller, L.H. (1986) Invasion of erythrocytes by malaria parasites: a cellular and molecular overview. *Annu. Rev. Microbiol.* 40, 451-477.
- 48 Bates, M.D., Newbold, C.I., Jarra, W. and Brown, K.N. (1988) Protective immunity to malaria: studies with cloned lines of *Plasmodium chabaudi chabaudi* in CBA/Ca mice. III. Protective and suppressive responses induced by immunization with purified antigens. *Parasite Immunol.* 10, 1-15.
- 49 Wood, J.C., Sales de Aguiar, J.C., Jarra, W., Ogun, S.A., Snounou, G. and Brown, K.N. (1989) In vivo selection of populations of *Plasmodium chabaudi chabaudi* AS resistant to a monoclonal antibody that reacts with the precursor to the major merozoite surface antigen. *Infect. Immun.* 57, 2128-2135.

Molecular and  
Elsevier

MOLBIO 014:

Si  
Schi.

Centre d

Molecular n  
characterization  
(Sm39) and me  
invertebrate de  
parasite and in  
The mollusc se  
species. The h  
could be speci  
the vesicles pr  
around sporoc

Key words: Se

## Introductio

Adaptatio  
developmen  
gies to eva  
molecular r  
of these po  
tosomes sh  
invertebrate  
non-specific  
as from the  
[3,4]. In thi  
port the im  
antigenic m

Corresponden  
logie et de Bi  
CNRS 624. In

Note: Nucleot  
been submitte  
number M 36

Abbreviations:  
SDS, sodium

0166-6851/90/

## Primary structure of the merozoite surface antigen 1 of *Plasmodium vivax* reveals sequences conserved between different *Plasmodium* species

(parasitic protozoa/malaria/vaccine/gene cloning)

HERNANDO A. DEL PORTILLO\*, SHIRLEY LONGACRE, ELISABETH KHOURI, AND PETER H. DAVID

Unité d'Immunoparasitologie, Institut Pasteur, 25 rue du Dr. Roux, Paris 75015, France

Communicated by William Trager, January 9, 1991 (received for review December 17, 1990)

**ABSTRACT** Merozoite surface antigen 1 (MSA1) of several species of plasmodia has been shown to be a promising candidate for a vaccine directed against the asexual blood stages of malaria. We report the cloning and characterization of the MSA1 gene of the human malaria parasite *Plasmodium vivax*. This gene, which we call *Pv200*, encodes a polypeptide of 1726 amino acids and displays features described for MSA1 genes of other species, such as signal peptide and anchoring sequences, conserved cysteine residues, number of potential N-glycosylation sites, and repeats consisting here of 23 glutamine residues in a row. When the nucleotide and deduced amino acid sequences of the MSA1 of *P. vivax* are compared to those of another human malaria parasite, *Plasmodium falciparum*, and to those of the rodent parasite *Plasmodium yoelii*, 10 regions of high amino acid similarity are observed despite the very different dG+dC contents of the corresponding genes. All of the interspecies conserved regions reside within the conserved or semiconserved blocks delimited by the sequences of different alleles of the MSA1 gene of *P. falciparum*.

The surface of the invasive merozoite of plasmodia constitutes one of the potential targets of a vaccine directed against the blood stages of malaria. Merozoite surface antigen 1 (MSA1), described by Holder and Freeman in 1982 (1), has been extensively studied in the human malarial parasite *Plasmodium falciparum* (reviewed in ref. 2). There are several allelic forms of this polymorphic high molecular weight antigen, and conserved, semiconserved, and variable regions can be found in the different alleles (3–5). The antigen is processed on the surface of the merozoite, although the exact stage at which processing occurs is subject to discussion (6). MSA1 has also been shown to bind in a specific manner to the surface of erythrocytes and could thus constitute one of the merozoite surface ligands involved in invasion of the erythrocyte (7).

A number of immunization experiments performed with parasite-derived or recombinant MSA1 or with MSA1 peptides in monkeys (reviewed in ref. 2) as well as in humans (8) point to this antigen as one of the most promising vaccine candidates against malaria asexual blood stages. *P. falciparum* is the only human malarial parasite for which the protective properties of the MSA1 have been assessed. Since protective immunity in malaria is species-specific (9), it is unlikely that a vaccine against one species will protect against others. Although *Plasmodium vivax* is the most widely distributed human malaria parasite, little is known about the properties of MSA1 in this species (10); this is partly due to the difficulty in obtaining large quantities of a parasite that cannot be maintained in continuous culture. The cloning and characterization of the gene coding for the MSA1 of *P. vivax*

should allow appropriate immunization studies to be performed with recombinant proteins.

A portion of the *P. vivax* MSA1 gene (Belem strain) has been previously characterized (11), and we present here the complete primary structure of this gene,† which we call *Pv200*. The organization of *Pv200* is similar to that of the MSA1 gene of *P. falciparum*, *Pf190* (3, 12), and to that of the rodent malaria parasite *Plasmodium yoelii*, *Py230* (13). There are 10 regions of high amino acid similarity conserved among the three parasite species. Since this molecule, like many other *P. vivax* antigens, is otherwise polymorphic (14, 15), such regions of interspecies conservation could be of importance in the development of an asexual stage malaria vaccine.

### MATERIALS AND METHODS

**Parasites.** The *P. vivax* Belem strain, adapted to *Saimiri* monkeys, was used for the production of DNA (11).

**Construction and Screening of Genomic DNA Libraries.** Two DNA libraries were constructed: (i) Library A. Genomic DNA was completely digested with *EcoRI* and 5 µg was fractionated on a 1% agarose gel. Fragments between 5 and 15 kilobases (kb) were electroeluted from a slice of the gel, extracted with phenol, and precipitated with ethanol. Pellets were washed, dried, and dissolved in double-distilled H<sub>2</sub>O. A 1-µg aliquot was ligated into the *EcoRI* arms of the  $\lambda$  vector *gtWES* (GIBCO/BRL) according to the supplier's instructions. The library was obtained by transforming LE392 competent cells and it was screened with a 1.9-kb DNA insert containing a portion of the *Pv200* gene, *Pv200/1.9* (see *Results*) (11).

(ii) Library B. A 0.5-µg sample of *HindIII*-digested DNA was ligated into the *HindIII* site of the vector pBR322 treated with calf intestinal alkaline phosphatase (Pharmacia) and the library was obtained by transformation of DH5  $\alpha$  competent cells. The library was screened with a 0.98-kb DNA insert corresponding to the first 0.98 kb from the 5' end of the *Pv200/1.9* clone (see *Results*).

All enzyme digestions and DNA manipulations were performed as recommended in Sambrook *et al.* (16).

**DNA Sequences.** Dideoxy chain termination sequences (17) were obtained by the production of exonuclease III overlapping deletion clones (18) or by the use of oligonucleotides (17-mers) synthesized on an Applied Biosystems PCR-Mate apparatus. Both DNA strands were sequenced for all the results presented here. Sequences were aligned and analyzed

Abbreviations: MSA1, merozoite surface antigen 1; ICB, interspecies conserved block; CB, conserved block.

\*Present address: Departamento de Parasitologia, Instituto de Ciencias Biológicas II-Universidade de São Paulo-Cidade Universitária, Avenida Professor Lineu Prestes 1374, São Paulo, CEP 05508, Brazil.

†The sequence reported in this paper has been deposited in the GenBank data base (accession no. M60807).

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.





reading frame of 1726 amino acids with a calculated molecular weight of 194,267 is contained within the genomic fragments. This is in agreement with the absence of introns in the genes coding for the MSA1 of other species.

There are a potential signal peptide and a hydrophobic membrane anchor sequence at residues 1–17 and 1710–1726, respectively. Furthermore, there are 12 potential N-glycosylation sites (Asn-Xaa-Thr/Ser) and 22 cysteines, 11 of which are located within the last 110 residues of the COO terminus of the molecule. The Pv200 sequence also contains a stretch of 23 glutamines at residues 726–748.

Comparison of the codon usage in the MSA1 genes of *P. vivax*, *P. falciparum*, and *P. yoelii* revealed that codons which have G or C in the third position are more frequent in *P. vivax*. Consequently, the dG+dC content of the Pv200 coding region is 43.4% and differs significantly from the dG+dC content of the coding regions of the MSA1 genes from *P. falciparum* (25.7%) and *P. yoelii* (31%).

**Comparisons of the Pv200, Pf190, and Py230 Sequences.** The deduced amino acid sequence from the Pv200 gene was computer-aligned with the sequences of the Pf190 (allele MAD20) (Fig. 2) and Py230 YM (Fig. 3) polypeptides. There is an overall identity of 35.6% and 34.3% with the *P. falciparum* and *P. yoelii* sequences, respectively.

Interestingly, 17 out of the 22 cysteines of the Pv200 polypeptide were located at similar positions with respect to the Pf190 and Py230 sequences. These similarities include the 11 and 10 cysteines found at the COO terminus of Pf190 and

Py230, respectively. In contrast, of 12 (Pv200), 15 (Pf190), and 11 (Py230) potential N-glycosylation sites, only 3 were conserved at the same positions between the *P. vivax* and the *P. falciparum* sequences, whereas only 1 was conserved between the *P. vivax* and *P. yoelii* sequences.

To determine the regions with an amino acid identity near 50% among the three parasite species, we combined the comparisons which had been made between Pv200–Pf190/Pv200–Py230 (this work) and Py230–Pf190 (13). Fig. 4 shows the result of such analysis. Seven ICBs were observed: ICB1, ICB2, ICB4, ICB5, ICB6, ICB8, and ICB10. Similarly, three other blocks (CB3, CB7, and CB9) were conserved between Pv200 and Pf190 but not between Pv200 and Py230 and thus could not be treated as bona fide ICBs. All these blocks reside within the conserved or semiconserved blocks of the Pf190 alleles (3).

## DISCUSSION

We report the complete primary structure of the MSA1 gene of the *P. vivax* Belem strain, Pv200. The general structure of the gene resembles that of the MSA1 genes described for *P. falciparum* and *P. yoelii*, with a number of homologous regions and other features such as (i) conserved cysteine residues at the COO-terminal region, (ii) number of potential N-glycosylation sites, and (iii) the presence of 23 glutamines in a row, which could correspond in *P. vivax* to the repeated sequences described in the MSA1 genes of other species.




FIG. 2. Comparison of the amino acid sequences of the *P. vivax* Belem strain Pv200 (upper sequence) and of the *P. falciparum* MAD20 Pf190 (lower sequence) (3). Sequences were aligned by using the program of Staden (19). Hyphens indicate gaps introduced for alignment; colons, identical residues; and periods, similar residues. Positions of the Pv200 cysteine residues conserved between these two proteins are denoted by ● and those that are not conserved, by ○. The position of the Pf190 blocks determined by the sequence of different alleles (3) is also shown; conserved blocks are indicated by unbroken underlines and overlines, semiconserved blocks are indicated by broken underlines and overlines, and variable blocks are unmarked.




FIG. 3. Comparison of the amino acid sequences of the *P. vivax* Belem strain Pv200 (upper sequence) and of the *P. yoelii* YM Py230 (lower sequence) (13). Sequences were aligned by using the program of Staden (19). Conventions are as in Fig. 2.

Malaria parasites have been divided evolutionarily into three groups according to the base composition of their DNA (22). One group, comprising avian, rodent, and falciparum malarias, presents a genome with a low dG+dC content (18%). Another, comprising the two monkey malarias *Plasmodium knowlesi* and *Plasmodium fragile*, presents a genome with a higher dG+dC content (30%). Finally, the group of *P. vivax* and *Plasmodium cynomolgi*, human and monkey malarias which cause relapses, has a genome presenting both low and high dG+dC components. This division implies that homologous genes and their proteins should be more similar within a group than between groups (22). Our observations show that in the case of the MSA1 genes and their proteins this prediction is supported only at the nucleotide level. Indeed, the low dG+dC content of the *Pf190* and *Py230* genes leads to a higher similarity, at the nucleotide level, between them than with *Pv200*. However, when the amino acid

composition is considered, *Pv200* and *Pf190* antigens show higher similarity and the overall distribution of their shared amino acids is more highly conserved than when *Pf190* and *Py230* are compared. That a higher amino acid similarity and closer overall distribution are observed in the *Pv200* and *Pf190* antigens despite their very different total dG+dC content most likely reflects the effects of positive selection within the human host. Accordingly, three regions of homology between the *Pv200* and *Pf190* antigens not conserved between the *Pv200* and the *Py230* antigens can be found (Fig. 4).

The analysis of the primary structure from different alleles of the MSA1 gene of *P. falciparum* allowed the definition of conserved, semiconserved, and variable regions within the molecule (3). One of the regions of amino acid identity higher than 45% conserved between the *Pf190* and *Py230* antigens resides within a variable block of one of the *Pf190* alleles and,




FIG. 4. Representation of the MSA1 antigen based upon amino acid conservation among the *Pv200*, *Pf190*, and *Py230* proteins (inner blocks) and upon *Pf190* alleles (outer blocks); solid-outline blocks, conserved areas; broken-outline blocks, semiconserved areas (7). Shaded boxes represent interspecies conserved blocks (ICBs) with greater than 48% identity among the three parasite species. Hatched boxes represent conserved blocks (CBs) with greater than 50% identity between *Pv200* and *Pf190* but not between *Pv200* and *Py230*. Open boxes represent areas of less than 45% identity. Positions of ICBs and CBs (amino acid residues of the *Pv200* sequence): ICB1, 1–50; ICB2, 107–200; ICB3, 274–319; ICB4, 348–387; ICB5, 620–691; ICB6, 796–895; ICB7, 1040–1088; ICB8, 1092–1153; ICB9, 1347–1464; and ICB10, 1622–1727.

consequently, Lewis (13) proposed the delimitation of new conserved blocks within the MSA1 antigen based on interspecies conservation. We decided to conduct a similar analysis; regions of 50 or more contiguous amino acids presenting 50% or higher identity among the three species [Pv200 vs. Pf190 and Pv200 vs. Py230 (this work) and Py230 vs. Pf190 (13)] are referred to as ICBs. Subsequently, the position of such ICBs with respect to the blocks delimited by sequences from different Pf190 alleles (3) was also examined.

All of the ICBs of MSA1 described here reside within the conserved or semiconserved blocks delimited by different alleles of the *P. falciparum* gene (Fig. 4). That such well-defined regions of MSA1 have been conserved among these three different malaria species could be explained because they are functionally or structurally important for the molecule, or because they are not immunogenic, or, finally, because immune responses against them do not block parasite growth (23). On the basis of these results, we predict that as sequences from other alleles of the *Pv200* gene are described, the general structure of the *Pv200* gene will comprise blocks that will be organized in a fashion similar to that of the blocks delimited by different *Pf190* alleles.

As for the protective properties of MSA1, most immunization trials with *P. falciparum* have used either the whole molecule or fragments from the NH<sub>2</sub>-terminal part (reviewed in ref. 2). In particular, the two peptides used in human vaccine trials belong to the regions we have defined as ICB1 and CB3 (8). This does not exclude other portions of MSA1; in particular, ICB10 corresponds to the most COO-terminal part of the molecule. The most remarkable aspect of this part of MSA1 is that it contains more than half of all the cysteine residues that are conserved in position among the three parasite species. Significantly, a protective monoclonal antibody against a discontinuous epitope of the *P. yoelii* MSA1 has been mapped to this region (24). Immunization trials with the MSA1 antigen of *P. vivax* have yet to be reported, and the potential protective properties of *Pv200* can only be extrapolated from experiments performed in other malarial species. The availability of the complete primary structure from the MSA1 gene of *P. vivax* should now allow the assessment of *Pv200* as a vaccine candidate.

We are grateful to Prof. Ruth Nussenzweig, Dr. Valerie Snewin, and Dr. Alexandra Levitt for critically reviewing the manuscript; to Ms. Christiane Duponchel for providing some of the oligonucleotides used for sequencing; to Dr. Michel Rabinovitch and Dr. Jurg Gysin for support; to Dr. Denise Mattei and Dr. Artur Scherf for helpful scientific discussions; and to Dr. Mario L. A. Bressan, Dr. Charlie Roth, and Dr. Michel Keller for computer assistance. This investigation received support from the United Nations Development Programme/World Bank/World Health Organization Special Pro-

gramme for Research and Training in Tropical Diseases (TDR) and from the Rockefeller Foundation.

- Holder, A. A. & Freeman, R. R. (1982) *J. Exp. Med.* **156**, 1528-1538.
- Holder, A. A. (1988) *Prog. Allergy* **41**, 72-97.
- Tanabe, K., Mackay, M., Goman, M., & Scaife, J. G. (1987) *J. Mol. Biol.* **196**, 273-287.
- Certa, U., Rotmann, D., Matile, H., & Reberliske, R. (1987) *EMBO J.* **6**, 4137-4142.
- Kimura, E., Mattei, D. M., di Santi, S. M., & Scherf, A. (1990) *Gene* **91**, 57-62.
- Pirson, P. J. & Perkins, M. E. (1985) *J. Immunol.* **134**, 1946-1951.
- Perkins, M. E. & Rocco, L. J. (1988) *J. Immunol.* **141**, 3190-3196.
- Patarroyo, M. E., Amador, R., Clavijo, P., Moreno, A., Guzman, F., Romero, P., Tascon, R., Franco, A., Murillo, L. A., Ponton, G., & Trujillo, G. (1988) *Nature (London)* **332**, 158-161.
- Becker, F. T. (1949) in *Malariaology*, ed. Boyd, M. F. (Saunders, Philadelphia), Vol. 2, pp. 1145-1157.
- David, P. H., Barnwell, J. W., & Mendis, K. N. (1990) in *New Generation Vaccines*, eds. Woodrow, G. C. & Levine, M. M. (Dekker, New York), pp. 531-544.
- del Portillo, H. A., Gysin, J., Mattei, D. M., Khouri, E., Udagama, P. V., Mendis, K. N., & David, P. H. (1988) *Exp. Parasitol.* **67**, 346-353.
- Holder, A. A., Lockyer, J., Odink, K. G., Sadhu, J. S., Riveros-Moreno, V., Nicholls, S. C., Hillman, Y., Davey, L. S., Tizard, M. L.-V., Schwart, R. T., & Freeman, R. R. (1985) *Nature (London)* **317**, 270-273.
- Lewis, A. P. (1989) *Mol. Biochem. Parasitol.* **36**, 271-282.
- Udagama, P. V., David, P. H., Peiris, J. S. M., Ariyaratne, Y. G., Perera, K. L. R. L., & Mendis, K. N. (1987) *Infect. Immun.* **55**, 2604-2611.
- Udagama, P. V., Gamage-Mendis, A. C., David, P. H., Peiris, J. S. M., Perera, K. L. R. L., Mendis, K. N., & Carter, R. (1989) *Am. J. Trop. Med. Hyg.* **42**, 104-110.
- Sambrook, J., Fritsch, E. F., & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab., Cold Spring Harbor, New York), 2nd Ed.
- Sanger, F., Nicklen, S., & Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 5463-5467.
- Henikoff, S. (1984) *Gene* **28**, 351-359.
- Staden, R. (1982) *Nucleic Acids Res.* **10**, 2951-2961.
- Saul, A. & Battistutta, D. (1990) *Mol. Biochem. Parasitol.* **42**, 55-62.
- Mackay, M., Goman, M., Bone, N., Hyde, J. E., Scaife, J., Certa, U., Stunnenberg, H., & Bujard, H. (1985) *EMBO J.* **4**, 3823-3829.
- McCutchan, T. F., Dame, J. B., Miller, L. H., & Barnwell, J. (1984) *Science* **225**, 808-811.
- Weber, J. L., Leininger, W. M., & Lyon, J. A. (1986) *Nucleic Acids Res.* **14**, 3311-3323.
- Burns, J. M., Jr., Majarian, W. R., Young, J. F., Daly, T. M., & Long, C. A. (1989) *J. Immunol.* **143**, 2670-2676.

Notice: This material may be  
protected by copyright law  
(Title 17 U.S. Code).

## Structure and expression of the gene for Pv200, a major blood-stage surface antigen of *Plasmodium vivax*

Helen L. Gibson<sup>1</sup>, Jeffrey E. Tucker<sup>1</sup>, David C. Kaslow<sup>2</sup>, Antoniana U. Krettli<sup>2</sup>,  
William E. Collins<sup>3</sup>, Michael C. Kiefer<sup>1</sup>, Ian C. Bathurst<sup>1</sup>, and Philip J. Barr<sup>1</sup>

<sup>1</sup>Chiron Corporation, Emeryville, CA, U.S.A.; <sup>2</sup>The Laboratory of Parasitic Diseases, National Institute of Allergy  
and Infectious Diseases, National Institutes of Health, Bethesda, MD, U.S.A.; and <sup>3</sup>Division of Parasitic Diseases,  
Center for Infectious Diseases, Centers for Disease Control, Public Health Service, U.S. Department of Health  
and Human Services, Atlanta, GA, U.S.A.

(Received 9 August 1991; accepted 21 September 1991)

Molecular cloning and structure analysis of the gene encoding the Pv200 protein of the Sal-1 strain of *Plasmodium vivax* revealed an overall identity of 34-37% when the deduced amino acid sequence was compared with the sequences of various major merozoite surface antigens of *Plasmodium falciparum*, *Plasmodium yoelii* and *Plasmodium chabaudi*. When the Sal-1 Pv200 sequence was compared with the corresponding sequence from the Belém strain of *P. vivax*, it was found that the two merozoite surface antigens were relatively well conserved with an overall amino acid sequence identity of 81%. A region of 23 repeated glutamine residues, found in the sequence of the Belém isolate was not found, however, in the Sal-1 sequence. Amino- and carboxy-terminal domains of the Pv200 protein were expressed in the yeast *Saccharomyces cerevisiae*. Each recombinant protein was shown to react with antibodies in sera from splenectomized Bolivian *Saimiri* monkeys that had been infected previously with *P. vivax*, and in human sera from individuals with a history of exposure to vivax malaria. The availability of recombinant DNA-derived Pv200 proteins will now allow a full assessment of their utility in the diagnosis and immunoprophylaxis of the benign tertian malaria associated with *P. vivax* infection.

Key words: *Plasmodium vivax*; Blood-stage antigen; Pv200; Malaria vaccine

### Introduction

Large scale in vitro parasite culture cannot provide sufficient quantities of either organisms or antigens to produce a malaria vaccine. Accordingly, peptide synthesis and recombinant DNA methodologies are being evaluated

extensively for the production of potential protective immunogens against the protozoan parasites that cause malaria. Because of the severity of the disease caused by *Plasmodium falciparum*, this species has represented the major target for such studies. Of the numerous potential subunit vaccine candidates from this organism, an antigen of special interest has been the major merozoite surface antigen Pf195. This antigen has been shown to reside on the surface of the schizont and, in a processed form, on the surface of the merozoite [1]. Vaccination studies with Pf195, isolated from cultured parasites, have led to high levels of protection against *P. falciparum* challenge in monkey model systems [2]. More recently, recombinant DNA-derived Pf195 antigens have been produced and studied as candidate vaccines against falciparum malaria [3-5].

Correspondence address: Philip J. Barr, Molecular Biology,  
Chiron Corporation, 4560 Horton Street, Emeryville, CA  
94608, USA.

Note: Nucleotide sequence data reported in this paper have  
been submitted to the GenBank™ data base with the accession  
number M75674.

Abbreviations: ADH2, alcohol dehydrogenase-2; CS, circum-  
sporozoite; GAPDH, glyceraldehyde-3-phosphate dehydrogen-  
ase; PAGE, polyacrylamide gel electrophoresis; PCR,  
polymerase chain reaction; TMD, transmembrane domain.

Although less virulent than *P. falciparum*, *Plasmodium vivax* is the causative agent of benign tertian fever, a form of malaria characterized by frequent and protracted relapses. Vaccination studies against this species have been less extensive than with *P. falciparum* and have been focused almost exclusively on the sporozoite lifecycle stage of the organism [6-9]. Recently, however, a polymorphic 200-kDa component of the *P. vivax* schizont surface was defined by monoclonal antibodies, and a partial genomic clone that encoded a portion of the antigen was isolated, structurally defined and expressed in bacteria [10]. Homology of this DNA fragment, and the more recently isolated full length gene sequence [11], with the Pf195 gene, together with immunolocalization studies using antisera to the expressed protein have suggested strongly that the Pv200 protein is functionally analogous to Pf195 [10]. Here we report the molecular cloning and structure analysis of the gene for Pv200 from the Sal-I strain [8] of *P. vivax*. We also demonstrate that amino- and carboxy-terminal domains of the protein produced in yeast can be used to detect antibodies both in monkeys and in humans previously infected with *P. vivax*.

## Materials and Methods

**Construction of *P. vivax* genomic DNA libraries.** Genomic libraries were prepared as follows. 500 ng of *P. vivax* genomic DNA, from the Sal-I strain, was digested with *Eco*RI and ligated into *Eco*RI digested  $\lambda$ ZAPII (Stratagene), packaged and introduced into *Escherichia coli* strain PLK-17. A similar *Hind*III digest, partially filled with dCTP and dTTP, was ligated into *Xba*I digested  $\lambda$ ZAPII that had been similarly filled with dATP and dGTP. The ligated DNA was packaged and transformed as above. Libraries of  $5 \times 10^7$  and  $4.1 \times 10^7$  independent clones were obtained, respectively.

**Screening of *P. vivax* DNA.** Two overlapping oligomers (45-mers), based on the sequence of

del Portillo et al. [10], were labeled by the oligomer primed extension method [12], hybridized in 40% formamide-containing buffer [13] at 37°C to a Southern blot of *Eco*RI-digested *P. vivax* DNA, and washed at 65°C in  $2 \times$  SSC/0.1% SDS. A 9.5-kb *Eco*RI fragment hybridized to this probe. Two overlapping oligomers (42- and 43-mers) based on the 5'-end of the *Eco*RI clone were used to probe a Southern blot of *Hind*III-digested *P. vivax* DNA in a similar fashion and hybridized to a 7.0-kb fragment. Similar hybridizations were carried out on library filters and plaque purified tertiary positives were excised using *E. coli* strain XL1-Blue (Stratagene) and plasmid DNA was retransformed into *E. coli* strain D1210 for further plasmid manipulations.

**Subcloning and DNA sequencing.** Plasmid DNA was isolated by the alkaline lysis method [13]. Overlapping restriction fragments were subcloned into M13 vectors and both strands were sequenced by the chain-termination method [14] using M13 primers as well as specific internal primers. DNA manipulations were essentially as described [13].

**Expression of amino- and carboxy-terminal domains of Pv200 in *Saccharomyces cerevisiae*.** The polymerase chain reaction (PCR) [15] was used to amplify DNA fragments from cloned Pv200 gene sequences. Appropriate restriction sites, and in-frame initiation and termination codons were incorporated into the PCR primers. Thus, for Pv200A, primers 5'-(dA-TGTCCCATGGAAACAGAAAGTTATAA-GCAG)-3' and 5'-(dCGCCCTCAACAAATCATAGTG)-3' were used to amplify an *Nco*I/*Eco*RI fragment from the *Hind*III clone 4B-3-9. A pBluescript polylinker primer, 5'-(dGTGGATCCCCCGGGCTGCAGG)-3' and the 3'-primer 5'-(dTCCAAGGTCGAC-TATGGATTTTGCAAATCACCAAATGT)-3' were used to amplify the contiguous *Eco*RI-*Sal*I fragment from the *Eco*RI clone 6.1-2. The PCR products were digested with the appropriate restriction enzymes and ligated into *Nco*I/*Sal*I digested pBS100 [7]. A *Bam*HI/*Sal*I fragment that contained the ADH2/GAPDH

hybrid promoter product was excised [17] for yeast expression amplified as an  $\lambda$  6.1-2 using the ATGGCAGAA7 and 5'-(dTTCAGAAAACTCC cloning into pBS24 above. pBS24-Pv200A and Pv200B were transformed into *Saccharomyces cerevisiae* derivative of ABY101 containing the *ADRI* gene [16], grown and induced in yeast cell lysate. Yeast cell lysates were analyzed by SDS-PAGE gels and by immunoblotting.

**Purification of Pv200A and Pv200B.** Cells that were harvested at 60 h in 1 M NaCl were washed with Triton X-100 and then with water. Pv200A was pelleted with 8 M urea. The solution was passed over a Mono Q ion exchange column (HCl/1 mM EDT) and was >95% pure by Coomassie Brilliant Blue staining. Yeast cells expressing Pv200B were harvested and lysed in a sir extract was then applied to a Fast Flow Q column. The immunoreactive fraction eluting at 0.15-0.20 M NaCl was collected, concentrated by gel filtration on Sepharose 6B in 4 M urea/10 mM  $\beta$ -mercaptoethanol.

**Serology.** Immunoblots were performed on monkey and human sera essentially as described [10]. Controls obtained from monkeys that had not been infected with *P. vivax* were used to assess specificity.

hybrid promoter [16] fused to the cloned PCR product was excised and cloned into pBS24 [17] for yeast expression. The Pv200B gene was amplified as an *NcoI/SalI* fragment from clone 6.1-2 using the primers 5'-(dCGGGAGCC-ATGGCAGAATCTGAGGCGCCTGAG)-3' and 5'-(dTTC CAAGGTCGACTAGCTAC-AGAAA CTCCCTCAAAGAG)-3'. Subcloning into pBS100 and pBS24 were as above. pBS24 plasmids containing the Pv200A and Pv200B genes were transformed into *Saccharomyces cerevisiae* strain JSC302, a derivative of AB116 [18] that overexpresses the *ADRI* gene [16]. Transformants were propagated and induced as described previously [7]. Yeast cell lysates were analyzed on 12.5% SDS-PAGE gels with Coomassie Blue staining, and by immunoblotting.

**Purification of recombinant proteins.** Yeast cells that were expressing Pv200A were harvested at 60 h and lysed with glass beads in 1 M NaCl/50 mM Tris/1 mM EDTA/0.1% Triton X-100 at pH 7.5 [7]. After centrifugation, Pv200A was extracted from the insoluble pellet with 8 M urea/100 mM Tris (pH 11.5). The solution was adjusted to pH 8.3 and passed over a Mono Q (Pharmacia) anion exchange column in 6 M urea/50 mM Tris-HCl/1 mM EDTA. Pv200A was found to be >95% pure by densitometric scanning of a Coomassie Blue-stained SDS-PAGE gel. Yeast cells expressing Pv200B were harvested and lysed in a similar manner. The 8 M urea extract was then adjusted to pH 9 and run over a Fast Flow Q (Pharmacia) ion exchange column. The immunoblot-positive peak eluting at 0.15–0.20 M NaCl in a 0–1 M gradient was collected, concentrated and separated by gel filtration on Superose 12 resin (Pharmacia) in 4 M urea/10 mM Tris/1 mM EDTA/10 mM  $\beta$ -mercaptoethanol at pH 8.

**Serology.** Immunoblots and ELISA assays on monkey and human sera were performed essentially as described [13]. Monkey sera were obtained from *Saimiri boliviensis* squirrel monkeys that had been used previously as controls to assess the efficacy of a recombinant

*P. vivax* circumsporozoite vaccine [8,9]. Anti-*Cynomolgus* conjugates were used to assay antibodies in monkey samples. Sera were taken 4 weeks after challenge with 10 000 *P. vivax* Sal-1 sporozoites. Human sera were collected from a population in Brazil, outside the endemic zone, that had encountered a single 50-day exposure to *P. vivax* [19]. Serum samples were taken 8 months after the outbreak, which was completely controlled by chemotherapy and insecticides. Enhancement of sensitivity in the human sera ELISAs was achieved using the 3, 3', 5, 5'-tetramethylbenzidine (TMB) as substrate and reagents as recommended by the manufacturer (Kirkegaard and Perry Laboratories Inc., Gaithersburg, MD).

## Results and Discussion

**Molecular cloning of the Pv200 gene.** Overlapping  $\lambda$ ZAPII phage clones were isolated that contained the large open reading frame encoding Pv200 (Fig. 1). Most of the Pv200 protein sequence is encoded by the *EcoRI* clone (6.1-2). The amino-terminal region is contained within the ca. 7-kb *HindIII* clone (4B-3-9). The full DNA sequence of a 5.83-kb composite *NdeI* fragment that includes the entire Pv200 coding sequence is shown (Fig. 2). Typical promoter elements [20] are found within the 5'-region of the sequence, and a consensus motif for efficient initiation of translation [21] is apparent around the proposed initiation codon.

**Structure of Pv200.** As deduced from the composite sequence, Pv200 is derived from a 1751-amino acid precursor protein that contains a typical amino-terminal secretory signal sequence [22], and a transmembrane domain (TMD) (Fig. 1). A predicted signal peptidase cleavage site [22] occurs after Cys19. Also, within the hydrophobic TMD (Fig. 2), is a stretch of 4 serine residues that is reminiscent of the signal for attachment of glycosylphosphatidyl inositol anchor sequences to proteins of various protozoans, including Pf195 [23].




Fig. 1. Restriction maps of the overlapping genomic DNA clones that encode Pv200, shown above a schematic representation of the encoded protein sequence. The flanking *NdeI* sites (Fig. 2, legend) are shown (N). Other sites used for mapping and DNA sequencing: *HindIII* (H), *EcoRI* (E), and *XbaI* (X). Potential asparagine-linked glycosylation sites are shown (♦), as is the serine/threonine-rich region (STR). The signal sequence (Sig) and putative transmembrane domain (TMD) are shown in black, and a potential proteolytic cleavage site (see text) is arrowed. Also shown schematically, are the regions Pv200A and Pv200B that were selected for expression in yeast.

Overall amino acid sequence identity between our Pv200 and Pf195 from various sources [24] is in the 36–37% range, although a region of more extensive identity (45%) exists in the amino-terminal region [10, and present study]. Similarly, amino acid sequence identities between Pv200 and the major merozoite surface antigens of *P. chabaudi chabaudi* and *P. yoelii yoelii* were 34.6% and 34.9%, respectively [25,26]. Also present is a potential proteolytic cleavage site identical to that defined previously for the generation of the carboxy-terminal p42 protein of the Pf195 precursor [27] (Figs. 1 and 2, arrowed). Also in common with Pf195, Pv200 does not contain multiple repetitive elements that are characteristic of many other proteins of malaria parasites. Six serine/threonine-rich motifs of 5 amino acids are noted, however (Figs. 1, 2). Also, the striking 23-glutamine residue repeat of the Belém strain Pv200 protein was not present in the sequence encoded by our cloned gene, but rather, was replaced by a 35-amino acid residue stretch that contained only 6

glutamine residues. This, together with several other smaller insertions and deletions accounts for the larger size of the Sal-1 Pv200 precursor over that of the 1726 amino acid Pv200 precursor from the Belém strain. Despite this size difference, the two Pv200 precursors are relatively well conserved, with an overall amino acid sequence identity of 81%. As with the major merozoite surface antigens from all *Plasmodium* species that have been studied thus far, this intra-species homology could be divided into areas of the protein of relatively

Fig. 2. Composite DNA sequence of a 5.83-kb *NdeI* fragment containing the Pv200 coding sequence. The 1751-amino acid large open reading frame is shown with proposed amino-terminal signal and carboxy-terminal membrane-spanning sequences underlined. Also underlined, are 13 asparagine residues that are potential sites for glycosylation. A 30-amino acid serine/threonine-rich region between amino acids 241 and 270 that contains 6 copies of the 5 amino acid repeat motif G.S.(S/T).(N S G).(S/T) is also noted (double arrow). A potential proteolytic cleavage site (see text) after amino acid Glu1356 is arrowed.

269  
-150 TCATCATACA  
1 N K A  
1 ATGAGGCGC  
51 G E N  
151 GGAGAAATG  
101 R E L  
301 AGAGATTGG  
151 E H L  
451 GAGCATCTAA  
201 L I I  
601 TTAATTATTG  
251 G S S  
751 GGTTCGCTCA  
301 E F F  
901 ATATTTTACA  
351 T T N  
1051 ACTACAATC  
401 A K H  
1201 GCAAGATGG  
451 G S D  
1351 GGATCTGATG  
501 K E E  
1501 AAAAAGAGT  
551 E E K  
1651 GAGGAGAAGA  
601 K L K  
1801 AAATTAAAGT  
651 V G H  
1951 GTCCAGATC  
701 M I A  
2101 ATGATGCCAG  
751 N A V  
2251 AATGCAGTAA  
801 A P A F  
2401 GCGCCAGCAGC  
851 K H I F  
2551 AAGCAGTTT  
901 S I Y D  
2701 TCCATATATG  
951 A Q S A  
2851 GCTCAGTCAGC  
1001 V G G N  
3001 GTCCAGGATG  
1051 T K V K  
3151 AGAATGCGAA  
1101 L T S L  
3301 CTCACGATTT  
1151 L K T Y  
3451 CTCAGTACTA  
1201 H I E L  
3601 AAGATCGAGT  
1251 Y Q E L  
3751 TACCAGAGTT  
1301 A G S V  
3901 GCAGGATCAGT  
1351 Q V T I  
4051 CAAGTAACAGC  
1401 T H I T  
4201 ACTACATATAC  
1451 K K L I  
4351 AAGAGTCTAT  
1501 D I K  
4501 GATGATATTA  
1551 V S K V  
4651 GTGAGCAAGGT  
1601 K E E V  
4801 AAAATGAAGT  
1651 H V P D  
4951 AATGTGCTGA  
1701 E C K H  
5101 GAATGTAAAT  
1751 L  
5251 CTTTAAAAATA  
5401 AAAAAAATA





low amino acid sequence identity separated by conserved or semi-conserved regions [11,24]. We have identified thirteen distinct regions of Pv200 that are typified by their amino acid sequence identity levels between the Sal-1 and Belém strains (Fig. 3). Although not fully comparable with the Pf195 variable blocks, or their junctions [11,24], the existence of this distinct pattern suggests the possibility that Pv200 diversity could be generated by intra-genic recombination of a limited number of alleles, as is the case with Pf195 [24,28]. Of further note, the carboxy-terminal region of the Sal-1 strain Pv200, equivalent to p42 of *P. falciparum*, is encoded to a large degree by 2 of the highly conserved gene segments (11 and 13) and exhibits 91% amino acid sequence identity with the corresponding region of the Belém strain protein.

**Expression of Pv200 in yeast.** Two domains of Pv200 were selected for expression studies (Fig. 1). The first, designated Pv200A, includes amino acids 20-462 of the Pv200 precursor. Pv200A represents, therefore, a protein of approx. 49 kDa from the amino-terminal

region of Pv200. A similarly expressed region of Pf195 can elicit good immunological responses against native Pf195 in mice and rabbits (these authors, S.P. Chang, G.S. Hui, unpublished observations). Furthermore, this region of Pf195, expressed in bacteria, has been shown previously to induce partial protection in *Aotus* monkeys that were subjected to blood stage challenge with *P. falciparum* [5]. The second domain that was expressed, Pv200B, containing amino acids 1357-1729 of the Pv200 precursor, is the homolog of the carboxy-terminal fragment of Pf195 that has been structurally defined as p42 [27]. This protein is of considerable importance since the carboxy-terminus of the major merozoite surface antigen has been implicated in the induction of a protective immune response against *P. falciparum* infection [26,29]. Additional studies on this *P. falciparum* antigen have indicated that secreted recombinant p42, from insect cells, is recognized by conformation-dependent antibodies [3].

Each gene construct was generated by PCR [15] and expressed in the yeast *S. cerevisiae*. Pv200A was produced at particularly high




Fig. 3. Schematic representation of amino acid sequence identity distributions between Sal-1 and Belém strain (11) Pv200 proteins. Shown here is the larger Sal-1 strain protein, with gaps scored as mismatches. The position of the putative proteolytic cleavage site for generation of the *P. vivax* p42 homologue is shown (arrowed).

Fig. 4. SDS-PAGE plasmids were lysed 0.2% bromophenol proteins visualized yeast cell lysates; 1: purified Pv200A a preta

levels (Fig. 4A, the insoluble fr detergent-conta be purified to > scheme. Pv200 levels (<20% buffer insoluble to greater than procedure. The

pressed region  
immunological  
in mice and  
g. G.S. Hui,  
Furthermore, this  
eria, has been  
al protection  
cted to blood  
um [5]. The  
sed. Pv200B,  
729 of the  
blog of the  
195 that has  
2 [27]. This  
nce since the  
r merozoite  
cated in the  
ine response  
26.29]. Addi-  
trum antigen  
nbinant p42.  
y conforma-

ated by PCR  
*S. cerevisiae*.  
cularly high




Fig. 4. SDS-PAGE analysis of recombinant Pv200 antigens. (A) Yeast cells transformed with control or recombinant plasmids were lysed in gel loading buffer containing 10% glycerol, 50 mM DTT, 3% SDS, 625 mM Tris-HCl pH 6.8, and 0.2% bromophenol blue. After 3 sequential boiling and freezing steps, the lysates were electrophoresed on 12.5% gels and proteins visualized by staining with Coomassie Blue. Lane 1, molecular weight markers (BioRad prestained); lane 2, control yeast cell lysates; lanes 3, 4, total lysates from yeast cells expressing Pv200A and Pv200B respectively. (B) Immunoblot of purified Pv200A and Pv200B using pooled human sera (see text for details). Lane 1, molecular weight markers (BioRad prestained); lane 2, control yeast cell lysate; lanes 3, 4, purified Pv200A and Pv200B, respectively.

levels (Fig. 4A, lane 3) and represented 81% of the insoluble fraction, after lysis in a nonionic detergent-containing buffer. This protein could be purified to >95% by a one step purification scheme. Pv200B was expressed at moderate levels (<20% of total yeast Triton X-100 buffer insoluble fraction), but could be purified to greater than 85% using a 2-step purification procedure. The purified Pv200A and Pv200B

proteins were shown to recognize antibodies in sera from individuals with a previous history of *P. vivax* infection [19]. Pooled human sera from such individuals, who were positive in their responses to the *P. vivax* CS protein (see below), was shown to further react with the recombinant merozoite surface antigens, by immunoblot analysis (Fig. 4B).

strain [11] Pv200  
of the putative

**Seroreactivity of recombinant Pv200 antigens.** Pv200A and Pv200B were used to analyze sera from *Saimiri* monkeys that were infected previously with the Sal-1 strain of *P. vivax* by sporozoite challenge [8,9]. Somewhat surprisingly, the ELISA titers from these animals were low and were not indicative of disease state. In only 2 animals did we observe high titer responses, and in each case this was against Pv200B (data not shown). The reasons for this absence of high titer responses are currently unknown. However, these relatively low overall titers may reflect the fact that this was the primary challenge of a group of monkeys that had not been exposed previously to *P. vivax* malaria. In contrast, we observed relatively high ELISA titers in a human population that had been subject to a single outbreak of vivax malaria (Fig. 5). ELISA titers against both Pv200A and Pv200B were above control values, as were titers against the recombinant CS protein (Vivax2) [30]. Noticeably, titers against Pv200B were consistently higher than those against Pv200A. The most direct explanation for this is that the carboxy-terminal region of Pv200 is simply more




Fig. 5. ELISA titers of human sera. Sera from individuals with overt infections, in which parasitemia was detected by thick blood smears, and who were treated with oral chloroquine (numbers 1–16). Three individuals were also included as controls, one positive (number 17) who had multiple *P. vivax* and *P. falciparum* infections, with titers of around 655000 and 24000 for Pv200A and B respectively, and 2 negative (numbers 18 and 19) who were never exposed to *P. vivax* infections. Individuals in each group were assayed for reactivity against the *P. vivax* CS protein using a recombinant CS protein (Vivax2) [30] as well as against Pv200 A and B.

immunogenic than the amino-terminal domain, at least in a primary infection in humans. Alternatively, Pv200B might possess greater conformational integrity than Pv200A when compared with the corresponding regions of the native proteins, and might thus be more antigenic in the ELISA format. A third explanation is that higher titers against Pv200B could reflect the more conserved structure of this region. For example, the region of the Sal-1 Pv200 protein defined by our Pv200A molecule shares 76% amino acid sequence identity with the corresponding region of the Belém strain Pv200 protein whereas, as mentioned above, the carboxy-terminal regions, corresponding to Pv200B, are 91% identical. In conclusion, we have demonstrated that recombinant Pv200 proteins produced in recombinant yeast are able to recognize antibodies in infected monkeys and humans. The production of these proteins in large quantities will allow further and more detailed studies of their antigenicity, and their potential use as diagnostic reagents. In addition, their ability to elicit protective immune responses in experimental animals can now be evaluated.

#### Acknowledgements

We thank our friends and collaborators for their encouragement and support. In particular, we thank Drs. Pablo Valenzuela and Lou Miller. We also thank Dr. Peter David for communication of results prior to publication, and Peter Anderson for preparation of the manuscript. This work was supported by Chiron Corporation and Ciba Geigy Ltd.

#### References

- Holder, A.A. and Freeman, R.R. (1982) Biosynthesis and processing of a *Plasmodium falciparum* schizont antigen recognized by immune serum and a monoclonal antibody. *J. Exp. Med.* 156, 1528–1538.
- Siddiqui, W.A., Tam, L.Q., Kramer, K.J., Hui, G.S.N., Case, S.E., Yamaga, K.M., Chang, S.P., Chan, E.B.T. and Kan, S.-C. (1987) Merozoite surface coat precursor protein completely protects *Aotus* monkeys against *Plasmodium falciparum* malaria. *Proc. Natl. Acad. Sci. USA* 84, 3014–3018.


- Murphy, V.F., A.A. (1990) Ex insect cells and secretion. *Paras*
- Müller, H.-M., Lombardi, S., Development of the major surf *falciparum*. *Infect*
- Herrera, S., He Caspers, P., Dö (1990) Immuniz *dium falciparum* *Proc. Natl. Acad*
- Arnot, D.E., Ba V., Nussenzweig, zoite protein of characterization *Science* 230, 815
- Barr, P.J., Gib Hollingdale, M.F sion in yeast of a use in a human 1160–1171.
- Collins, W.E., N bush, T.K., Nard Young, J.F., Wa: H.L., Barr, P.J. Broderick, J.R., V.K. and Wilson. *sciurus boliviensis* based on the circ *vivax*. *Am. J. Tro*
- Collins, W.E., Bathurst, I.C., N G.H., Barr, P.J., E V.K., Stanfill, P. (1990) Further stu *sciurus boliviensis* the circumsporozo *J. Trop. Med. Hy*
- del Portillo, H.A., Udagama, P.V., *Plasmodium vivax* blood-stage surface 353.
- del Portillo, H.A., P.H. (1991) Prima antigen 1 of *Pl* conserved between *Natl. Acad. Sci. U*
- Feinberg, A. and V radiolabeling DNA to high specific act
- Sambrook, J., Ma Molecular Cloning Cold Spring Harb Harbor, NY.
- Sanger, F., Nicklen sequencing with c *Natl. Acad. Sci. U*
- Saiki, R.K., Scharf, G.T., Erlich, H.A. amplification of , restriction site an; anemia. *Science* 230

terminal do-  
infection in  
might possess  
than Pv200A  
responding re-  
might thus be  
rmat. A third  
against Pv200B  
d structure of  
ion of the Sal-  
our Pv200A  
acid sequence  
region of the  
areas, as men-  
ninal regions.  
% identical. In  
d that recom-  
d in recombi-  
antibodies in  
the production  
ties will allow  
dies of their  
l use as diag-  
neir ability to  
uses in experi-  
uated.

laborators for  
rt. In particu-  
zuela and Lou-  
ter David for  
to publication.  
aration of the  
supported by  
Geigy Ltd.


(1982) Biosynthesis  
*falciparum* schizont  
and a monoclonal  
338.  
K.J., Hui, G.S.N.,  
S.P., Chan, E.B.T.  
face coat precursor  
monkeys against  
c. Natl. Acad. Sci.

- 3 Murphy, V.F., Rowan, W.C., Page, M.J. and Holder, A.A. (1990) Expression of hybrid malaria antigens in insect cells and their engineering for correct folding and secretion. *Parasitology* 100, 177-183.
- 4 Müller, H.-M., Früh, K., von Brunn, A., Esposito, F., Lombardi, S., Crisanti, A. and Bujard, H. (1989) Development of the human immune response against the major surface protein (gp190) of *Plasmodium falciparum*. *Infect. Immun.* 57, 3765-3769.
- 5 Herrera, S., Herrera, M.A., Perlaza, B.L., Burki, Y., Caspers, P., Döbeli, H., Rotmann, D. and Certa, U. (1990) Immunization of Aotus monkeys with *Plasmodium falciparum* blood-stage recombinant proteins. *Proc. Natl. Acad. Sci. USA* 87, 4017-4021.
- 6 Arnot, D.E., Barnwell, J.W., Tam, J.P., Nussenzweig, V., Nussenzweig, R.S. and Enea, V. (1985) Circumsporozoite protein of *Plasmodium vivax*: gene cloning and characterization of the immunodominant epitope. *Science* 230, 815-818.
- 7 Barr, P.J., Gibson, H.L., Enea, V., Arnot, D.E., Hollingdale, M.R. and Nussenzweig, V. (1987) Expression in yeast of a *Plasmodium vivax* antigen of potential use in a human malaria vaccine. *J. Exp. Med.* 165, 1160-1171.
- 8 Collins, W.E., Nussenzweig, R.S., Ballou, W.R., Ruebush, T.K., Nardin, E.H., Chulay, J.D., Majarian, J.F., Young, J.F., Wasserman, G.F., Bathurst, I.C., Gibson, H.L., Barr, P.J., Hoffman, S.L., Wasserman, S.S., Broderick, J.R., Skinner, J.C., Procell, P.M., Filipinski, V.K. and Wilson, C.L. (1989) Immunization of *Saimiri sciureus boliviensis* monkeys with recombinant vaccines based on the circumsporozoite protein of *Plasmodium vivax*. *Am. J. Trop. Med. Hyg.* 40, 455-464.
- 9 Collins, W.E., Nussenzweig, R.S., Ruebush, T.K., Bathurst, I.C., Nardin, E.H., Gibson, H.L., Campbell, G.H., Barr, P.J., Broderick, J.R., Skinner, J.C., Filipinski, V.K., Stanfill, P.S., Roberts, J.M. and Wilson, C.L. (1990) Further studies on the immunization of *Saimiri sciureus boliviensis* with recombinant vaccines based on the circumsporozoite protein of *Plasmodium vivax*. *Am. J. Trop. Med. Hyg.* 43, 576-583.
- 10 del Portillo, H.A., Gysin, J., Mattei, D.M., Khouri, E., Udagama, P.V., Mendis, K.N. and David, P.H. (1988) *Plasmodium vivax*: cloning and expression of a major blood-stage surface antigen. *Exp. Parasitol.* 67, 346-353.
- 11 del Portillo, H.A., Longacre, S., Khouri, E. and David, P.H. (1991) Primary structure of the merozoite surface antigen 1 of *Plasmodium vivax* reveals sequences conserved between different *Plasmodium* species. *Proc. Natl. Acad. Sci. USA* 88, 4030-4034.
- 12 Feinberg, A. and Volgelstein, B. (1984) A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* 137, 266-267.
- 13 Sambrook, J., Maniatis, T. and Fritsch, E.F. (1989) *Molecular Cloning. A Laboratory Manual*, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- 14 Sanger, F., Nicklen, S. and Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
- 15 Saiki, R.K., Scharf, S., Faloona, F., Mullis, K.B., Horn, G.T., Erlich, H.A. and Arnheim, N. (1985) Enzymatic amplification of  $\beta$  globin genome sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 230, 1350-1354.
- 16 Shuster, J.R. (1989) Regulated transcriptional systems for the production of proteins in yeast: regulation by carbon source. In: *Yeast Genetic Engineering* (Barr, P.J., Brake, A.J. and Valenzuela, P., eds.), pp. 83-108. Butterworth, Stoneham, MA.
- 17 Sabin, E.A., Lee-Ng, C.T., Shuster, J.R. and Barr, P.J. (1989) High-level expression and in vivo processing of chimeric ubiquitin fusion proteins in *Saccharomyces cerevisiae*. *Bio/Technology* 7, 705-709.
- 18 Barr, P.J., Cousens, L.S., Lee-Ng, C.T., Medina-Selby, A., Masiarz, F.R., Hallewell, R.A., Chamberlain, S.H., Bradley, J.D., Lee, D., Steimer, K.S., Poulter, L., Burlingame, A.L., Esch, F. and Baird, A. (1988) Expression and processing of biologically active fibroblast growth factors in the yeast *Saccharomyces cerevisiae*. *J. Biol. Chem.* 263, 16471-16478.
- 19 Fontes, C.J., Bathurst, I. and Kretzli, A.U. (1991) *Plasmodium vivax* sporozoite antibodies in individuals exposed during a single malaria outbreak in a nonendemic area. *Am. J. Trop. Med. Hyg.* 44, 28-33.
- 20 Lewis, A.P. (1990) Sequence analysis upstream of the gene encoding the precursor to the major merozoite surface antigens of *Plasmodium yoelii*. *Mol. Biochem. Parasitol.* 39, 285-288.
- 21 Kozak, M. (1984) Compilation and analysis of sequences upstream from the translational start site in eukaryotic mRNAs. *Nucleic Acids Res.* 12, 857-872.
- 22 von Heijne, G. (1986) A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* 14, 4683-4690.
- 23 Smythe, J.A., Coppel, R.L., Brown, G.V., Ramasamy, R., Kemp, D.J. and Anders, R.F. (1988) Identification of two integral membrane proteins of *Plasmodium falciparum*. *Proc. Natl. Acad. Sci. USA* 85, 5195-5199.
- 24 Tanabe, K., MacKay, M., Goman, M. and Scaife, J.G. (1987) Allelic dimorphism in a surface antigen gene of the malaria parasite *Plasmodium falciparum*. *J. Mol. Biol.* 195, 273-287.
- 25 Deleersnijder, W., Hendrix, D., Bendahman, N., Hanegreefs, J., Brijs, L., Hamers-Casterman, C. and Hamers, R. (1990) Molecular cloning and sequence analysis of the gene encoding the major merozoite surface antigen of *Plasmodium chabaudi chabaudi* IP-PC1. *Mol. Biochem. Parasitol.* 43, 231-244.
- 26 Holder, A.A. and Freeman, R.R. (1984) Characterisation of a high molecular weight protective antigen of *Plasmodium yoelii*. *Parasitol.* 88, 211-219.
- 27 Heidrich, H.-G. (1988) Isolation and functional characterization of *Plasmodium falciparum* merozoite antigens. *Biol. Cell* 64, 205-214.
- 28 Peterson, M.G., Coppel, R.L., Moloney, M.B. and Kemp, D.J. (1988) Third form of the precursor to the major merozoite surface antigens of *Plasmodium falciparum*. *Mol. Cell. Biol.* 8, 2664-2667.
- 29 Burns, J.M., Jr., Daly, T.M., Vaidya, A.B. and Long, C.A. (1988) The 3' portion of the gene for a *Plasmodium yoelii* merozoite surface antigen encodes the epitope recognized by a protective monoclonal antibody. *Proc. Natl. Acad. Sci. USA* 85, 602-606.
- 30 Bathurst, I.C., Gibson, H.L., Kansopon, J., Hahm, B.K., Hollingdale, M.C. and Barr, P.J. (In Press) Muramyl peptide adjuvants for *Plasmodium falciparum* and *P. vivax* circumsporozoite vaccines in rodent model systems. *Biotech. Ther.*




~~Fig. 2~~

~~A~~ Fig. 2A



~~B~~ Fig. 2B






Fig. 3e Fig. 3c - Fig. 3x  
5/16

DNA sequence of the native (gp190n) and of the synthetic gene (gp190s) for gp190 from FCG-1

```
AS      M K I I F F L C S F L F F I I N T Q C V T H E S Y Q E      27
gp190n      G A T T A T T A A A T A A T A A T A G T A A
gp190s CCGACGGGATGAAATCATTTCTCTCTCTGTTCAATTCCTGTTTTTATCATCACTACTGCGGTGACCCAGCAATCCTATCAGGAG      90
Hlu I

AS      L V K K L E A L E D A V L T G Y S L F O K E K H V L N E G T      57
gp190n      T C A A A A A T G A T T T T A T A A A A T A A A
gp190s CTGGTTAAGAAACTGGAAGCTTTGGAAGATGCGGTCCTTACCGGATACAGCCTGTTCCAGAAGGAGAAGATGCTGCTGAATGAAGGGACG      180

AS      S G T A V T T S T P G S K G S V A S G G S G G S V A S G G S      87
gp190n      A A T T T T A G T A T T C A T A C A T T A T C A
gp190s AGTGGCAGCGGCTTACAACGACACACCGGTTCTAAAGGGTCTGTGGCTACCGGTGGCTCCGGTGGGTCTGTGGCCTCTGGGGGTTCC      270

AS      V A S G G S V A S G G S V A S G G S G N S R R T N P S D N S      117
gp190n      T T A T T C A T T T T T C A T A T T T C A A C T A T A T T A
gp190s GTCCCTCTCGGCGGCGGCTGCGCATCAGGTGGCTCAGTGGCAAGCGCGGTTCCGGGAACAGTGAAGAACCAATCCTCTGACAACCTCT      360

AS      S D S D A K S Y A D L K H R V R N Y L L T I K E L K Y P O L      147
gp190n      T A T T A T T A A A C T C T G T A A A C A T T T A C C
gp190s AGCGATTTCGCGGAGCTCTACGCGGAGCTCAAGCAGCGAGTGAGAACTATCTCTCACTATCAAGGAGCTGAAGTACCCACAGTTG      450

AS      F D L T N H M L T L C D N I H G F K Y L I D G Y E E I N E L      177
gp190n      T T A T A T T T T T T T T T T T T A T A T A T A T A
gp190s TTCCACCTCACTAATCATATGCTGACACTGTGTGATAACATTCTGGCTTCAAATATCTGATTGACGGTTACGAAGAGATCAATGAACTC      540

AS      L Y K L N F Y F D L L R A K L N D V C A N D Y C Q I P F N L      207
gp190n      T A T A A C T T T T A T A A T A T A T T T T T T C T
gp190s CTGTACAAGTTGAATTTCTACTTCGACTTGGTAAGGGCCAAACTGAATGACGTTTCCGCCAATGACTATTGTCAAATTCCTCAATTTG      630

AS      K I R A N E L D V L K K L V F G Y R K P L D N I K D N V G K      237
gp190n      A T C T A T A A C T A A C T G A A A A T A T T A T A
gp190s AAGATCAGAGCCCAACGAGTTGGACGCTATGAAGAAGTTGGTCTTCGGATATCCCAAGCCTCTCGACAACATCAAGGACATGTGGGAAAG      720

AS      H E D Y I K K N K K T I E N I N E L I E E S K K T I D K N K      267
gp190n      C A A A A T A T A T A T A T A T A T A T A T A T
gp190s ATGGAAGATTATATTAATAAAGATAAAGAAGACCATCGAGAACATTACGAGCTGATCGAAGAATCCAAAAGACCATAGACAAAAATAAG      810

AS      N A T K E E E K K K L Y Q A Q Y D L S I Y N K Q L E E A B N      297
gp190n      T A A A A A A A T A T T T T T T C T A T A A T
gp190s AATGCAACCAAGGAGGAAGAAAAGAAGAGTTGTACCGAGCCCAAGTACGACCTGTCCATCTATAACAAACAGCTTGAAGAAGCCCATAC      900

AS      L I S V L E K R I D T L K K N E N I K E L L D K I N E I K N      327
gp190n      T A A T T A A A T T T A A A C T G T A T T A A
gp190s CTGATCAGCTACTGGGAAGCGCATAGACACCTCAAGAAGAATGAAAAATATCAAGAAGCTGCTCGACAAGATTAATGAAATTAAGAAT      990

AS      P P P A N S G N T P N T L L D K N K K I E E H E K E I K E I      357
gp190n      C A G T A T A A T T C T T A A C A A A A A A A T
gp190s CCTCGCCAGCAACTCTGGGAACACCCCTAACAGCGCTGCTGGACAAGAACAAGAAGATAGAGGAGCAGGAGAAGAGATCAAGAGATC      1080

AS      A K T I K F N I D S L F T D P L E L E Y Y L R E K N K N I D      387
gp190n      T A T T T A G T A A A T A A T A A T A A T A
gp190s GCCAAAACCATTAAGTTCAACATAGATTCTCTTTTACTGATCCCTCTGAGCTGGAGTACTACTTGAGAGAGAAGAATAAGAATATAGAC      1170


AS      I S A K V E T K E S T E P N E Y P N G V T Y P L S Y N D I N      417
gp190n      A A G T A G T A T C A A A A A A A A T T T T A
gp190s ATCTCGCCCAAGTGGAGACAAAGGAATCAACCGAACCTAATGAATATCCCAATGGTGTGAGCTACCTCTGTCTTATAACGATATCAAC      1260

AS      N A L N E L N S F G D L I N P F D Y T K E P S K N I Y T D N      447
gp190n      T C T A T A T T C T T T A T A T A T A A A G A C A T T
gp190s AACGCTCTCAACGAGCTCAATAGCTTCGGTGACTTGATTAACCCCTTCGATTATACGAAGAAGCCTCTGAAGAATATCTACACAGACAAT      1350

AS      E R K K F I N E I K E K I K I E K K K I E S D K K S Y E D R      477
gp190n      A A A C A T T A A T A A A A A A T C T A T C A A
gp190s GAGAGAAGAAGTTTATCAACGAAATCAAGGAGAAGATCAAAATTCAGAAGAAGAAATTCAGAGTGAAGAAGAAAGTTACGAAGACCCG      1440

AS      S K S L N D I T K E Y E K L L N E I Y D S K F N N N I D L T      507
gp190n      T C T G T C T A A A A A A A T A T A T A T A T A
gp190s AGCAAAAGTCTAAACGATATCACTAAAGAGTATGAAAAGCTGTGAACGAGATCTATGATTCCAAATTCACAAATACATCGACCTGACC      1530

AS      N F E K H M G K R Y S Y K V E K L T H H N T F A S Y E N S K      537
gp190n      T A T A A T A T T T T T T T T T T T T T A
gp190s AACTTCGAGAAAATGATGGGAAAACGCTACTCTTACAAGTGGGAGAACTGACACACCATAATACCTTTGCTATCCTATGAGAATCTAAG      1620
```



6/16

AS H N L E K L T K A L K Y M E D Y S L R N I V V E K E L K Y Y 567  
gp190<sup>n</sup> A T A A A A T T A A T A A T A T A T  
gp190<sup>s</sup> CATAATCTTGAGAGCTCACCAGGCTCTTAAGTATATGGAGGACTATTCTCTCGGAAACATTGTTGTGGAGAAAGAAGCTAAAGTATTAC 1710

AS K N L I S K I E N E I E T L V E N I K K D E E Q L F E K K I 597  
gp190<sup>n</sup> A T A C A A T T A A T A T A A A C T A A A  
gp190<sup>s</sup> AAGAATCTCATAAGTAAGATCGAAAACGAGATCGAGAGCGCTTGTGGAACATTAGAAGGATGAAGAAGCTGTTGTTGAGAAGAAGATT 1800

AS T K D E N K P D E K I L E V S D I V K V Q V Q K V L L M N K 627  
gp190<sup>n</sup> T A A T T A A A T C A A T A A T T A T A A  
gp190<sup>s</sup> ACAAAGACGAAAATAAACAGATGAGAAGATCTCGGAGGCTCTCGATATTGTTAAAGTCCAAGTGCAGAGGCTGCTCTCATGAACAAG 1890

AS I D E L K K T Q L I L K N V E L K B N I H V P N S Y K Q E N 657  
gp190<sup>n</sup> C T A A T G T A A T A A T C T C C A A A  
gp190<sup>s</sup> ATTGATGAAGTCAAGAAGACTCAACTCATCTGGAAGAACCTGGAGTTAAACATAATATACATGTCGGAATAGTTATAAGCAGGAGAAT 1980

AS K Q E P Y Y L I V L K K E I D K L K V F M P K V E S L I N E 687  
gp190<sup>n</sup> A T T T A T G T G A A T T T A T G A A T C A T A T  
gp190<sup>s</sup> AAGCAGGAACCATACTACCTCATCTGACTCAAGAAAGAGATAGACAACTGAAAGTGTTCATGCCCAAGTGCAGAGGCTGATCAACGAA 2070

AS E K K N I K T E G Q S D N S E P S T E G E I T G Q A T T K P 717  
gp190<sup>n</sup> A A A A A T A G T G A A A C A A A A A A A A A A  
gp190<sup>s</sup> GAGAAGAAAGCAATTAAGTCAAGGACAGTCAAGTAAGTCCGAGCTTCACAGAGGAGAGATAACGGGACAGGCTACCAACAGGCCC 2160

AS G Q Q A G S A L E G D S V Q A Q A Q E Q K Q A Q P P V P V P 747  
gp190<sup>n</sup> A A A T T A A T C A A A A A A A A A A A A A A  
gp190<sup>s</sup> GGACAACAGGCGCGTTCAGCTCTCGAAGGCGATAGCTGCAAGCTCAAGCACAAGAGCAGAAGCAGGCAAGCTCCAGTCCAGTCCG 2250

AS V P E A K A Q V P T P P A P V N N K T E N V S K L D Y L E K 777  
gp190<sup>n</sup> A A A A C A A A A A A A A A A A A A A A A  
gp190<sup>s</sup> GTTCAGAGGCTAAAGTCAAGTCCCTACACACAGCTCCTGTGAATAACAGAGCAGAGATGTCAGCAAGCTGGACTACCTTGAGAAG 2340

AS L Y E F L N T S Y I C H K Y I L V S E S T M N E K I L K Q Y 807  
gp190<sup>n</sup> T A A T T A T A T A T T G T A T C A A A T A A T  
gp190<sup>s</sup> CTCTATGAGTTCCTGAATACATCTACATCTGCCACAAATATATCTCTGCTCTCACAGCACTATGAACGAGAAGATTCTTAAACAGTAC 2430

AS K I T K E E S K L S S C D P L D L L F N I Q N N I P V M Y 837  
gp190<sup>n</sup> A T A G A A C T A A G T A T A T A T A A T A A T  
gp190<sup>s</sup> AAGATAACCAAGGAAGAGGAGAGTAACTGTCTCTGTGATCCACTGGAGCTGCTGTTCAATATCCAGAACACATTCCCGTTATGTAT 2520

AS S M F D S L N N S L S Q L F M E I Y E K E M V C N L Y K L K 867  
gp190<sup>n</sup> T T A A G T A A A T A T A A A A T T T A T G  
gp190<sup>s</sup> TCTATGTTGATAGCCTCAACAATCTCTCTCTCAACTGTTTCATGAGATATATGAGAAGGAGATGGTCTGCAAGCTGTATAAACTCAAA 2610

AS D N D K I K N L L E E A K K V S T S V K T L S S S S M Q P L 897  
gp190<sup>n</sup> T T A A T T A T A G A A A A A A A A A A A A  
gp190<sup>s</sup> GACAACGACAGATTAAGAAGCTTCTGAGGAAGCTAAGAAGTCTCCAGCTCTGTAAAACCTCTCTCTCCAGCTCCATGCAACCACTG 2700

AS S L T P Q D K P E V S A N D O T S E S T N L N N S L K L F E 927  
gp190<sup>n</sup> A T A G T A A T A T A T T T A A T T A T T G T A G T A T A  
gp190<sup>s</sup> TCTCTCACAGCTCAAGACAAGCCGAGTGAAGGCTAAGCAGCAGCTCTCACTGACCAACCTTAATAACTCACTGAAAGCTGTTTGAG 2790

AS N I L S L G K N K N I Y Q E L I G O K S S E N F Y E K I L K 957  
gp190<sup>n</sup> A T A G T A A C A T A T A A T A A G T A T A T A T A  
gp190<sup>s</sup> AACATCTGTCTCTCGCAAGAATAAGAACATCTACCAAGAAGTATGGAAGCAAAATCTGCGAGAAGTCTACGAGAAGATAGTCAA 2880

AS D S D T F Y N E S F T N F V K S K A D D I N S L N D E S K R 987  
gp190<sup>n</sup> T T T T T A T C T T A T T A T T T A T T A T G T A A G  
gp190<sup>s</sup> GACAGCGACACATCTATAACGAGAGCTTCACTAAGTCTGTAATCTAAAGCCGATGATCAACTCTCTTAACGATGAATCTAAAGCT 2970

AS K K L E E D I N K L K K T L O L S F D L Y N K Y K L K L E R 1017  
gp190<sup>n</sup> A T A A T T A T A A A T T A G T A T C T T A T T A T A T A  
gp190<sup>s</sup> AAGAAGCTGGAAGAGGACATCAATAAGCTGAAGAAGCACTGCAACTGAGCTTGCAGCTGTACAACAGTACAACTGAAAGCTGGAGAGA 3060

AS L F D K K K T V G K Y K M Q I K K L T L L K E O L E S K L N 1047  
gp190<sup>n</sup> T A T T A A A T T A A A A T A A C T T A T A A A A T A T C A T G T  
gp190<sup>s</sup> CTCTTCGACAAGAAGACAGCTCGGCAAGTATAAGATGCAGATCAAGAAGTTGACTTCTGCTCAAGGAGCAGCTTGAAGCAAACTCAAC 3150

AS S L N N P K H V L O N F S V F F N K K K E A E I A E T E N T 1077  
gp190<sup>n</sup> T T C A G T T A A T T T T A A A A T A A A A T A  
gp190<sup>s</sup> TCAGTCAACATCCGAACAGTACTGCGAGAAGTCTCAGTGTCTCTCAACAAGAAGAAGGAGGAGATCGCGGAGACAGAGAACCT 3240

AS L E N T K I L L K H Y K G L V K Y Y N G E S S P L K T L S E 1107  
gp190<sup>n</sup> T A A A A A T A T G T T A T T A T A A A A T A A T A A T A A G T A  
gp190<sup>s</sup> CTGGAGAACACCAAGATCTCTCTCAACACTACAAAGGCTCGTCAAGTATTATAATGGGAGTCTCTCTCTGGAAGACTCTCTCCGAG 3330

AS E S I Q T E D N Y A S L E N F K V L S K L E G K L K D N L N 1137  
gp190<sup>n</sup> A T C A T A A A T T T A A A T A A T A A G A T A A A T A T T A T  
gp190<sup>s</sup> GAGAGCATCCAGACCGAGGATAACTAGCCAGGCTCGAGAAGTTCAGGCTCTGTGTAAAGCTCGAAGGCAAGCTGAAGGACAACCTGAAC 3420

7 / 16



~~Fig. 3D~~  
Fig. 3Y



8 / 16



	N'-terminus	C'-terminus
gp190s1 Sequence		
DNA Sequence	GC <u>ACGCGTATGAAAATC</u> ----- AGCTCTAATTAAATAGGCGGGCCGCATCGATGGC	
AA Sequence	Mlu I Met Lys Ile ----- Ser Ser Asn stop codon Not I Cla I	
AA Position	1 2 3 ----- 1619 1620 1621	
gp190s2 Sequence		
DNA Sequence	GC <u>GGATCCGTGACCCAC</u> ----- AGCTCTAATTAAATAGGCGGGCCGCATCGATGGC	
AA Sequence	BamH I Val Thr His ----- Ser Ser Asn stop codon Not I Cla I	
AA Position	20 21 22 ----- 1619 1620 1621	

~~Fig. 5C~~  
Fig. 5B

13/16





16/16

3

2

1

~~Fig. 6c~~ Fig. 6b


190

108

89


77

61



REPLACEMENT DRAWING  
U.S. Patent Application No. 09/269,874  
Edward J. Baba (Reg. No. 52,581)  
(650) 833-7731

FIG. 1







FIG. 2A









FIG. 2B









FIG. 3B










FIG. 3D

AS	S G T A V T T S T P G S K G S	72
gp190n	A A T T T A G T A	
gp190s	AGTGGCACGGCCGTTACAACCAGCACACCCGGTTCCTAAAGGTCT	225
AS	V A S G G S G G S V A S G G S	87
gp190n	T TCA T A C A T T A T C A	
gp190s	GTGGCTAGCGGTGGCTCCGGTGGGTCTGTGGCCTCTGGGGGTCC	270
AS	V A S G G S V A S G G S V A S	102
gp190n	T T A T TCA T T T TTCA	
gp190s	GTCGCCTCCGGCGGCAGCGTGGCATCAGGTGGCTCAGTGGCAAGC	315
AS	G G S G N S R R T N P S D N S	117
gp190n	T A T TTCAA C T A T A T T A	
gp190s	GGCGGTTCCGGGAACAGTCTGAAGAACCAATCCATCTGACAACTCT	360
AS	S D S D A K S Y A D L K H R V	132
gp190n	T A T T A T T T A A A A	
gp190s	AGCGATTCCGACGCCAAGTCTCTACGCCGACCTCAAGCACCAGTG	405
AS	R N Y L L T I K E L K Y P Q L	147
gp190n	C T CT GT A A A C A T T AC C	
gp190s	AGAAACTATCTCTCACTATCAAGGAGCTGAAGTACCCACAGTTG	450



FIG. 3E

AS	F	D	L	T	N	H	M	L	T	L	C	D	N	I	H	162	
gp190n	T	T	A					T	A	T	T						
gp190s	T	T	G	A	C	C	T	C	A	T	A	T	G	C	T	G	495
AS	G	F	K	Y	L	I	D	G	Y	E	E	I	N	E	L	177	
gp190n	T				T	A		T	A	T	A	T					
gp190s	G	G	T	T	C	A	A	A	T	A	T	G	A	C	C	T	540
AS	L	Y	K	L	N	F	Y	F	D	L	L	R	A	K	L	192	
gp190n	T	A	T	A	A	C	T	T	T	T	A	T	A	A	T		
gp190s	C	T	G	T	A	C	A	A	G	T	T	C	T	A	C	T	585
AS	N	D	V	C	A	N	D	Y	C	Q	I	P	F	N	L	207	
gp190n	T	A	T	T							A	T					
gp190s	A	A	T	G	A	C	C	C	A	A	T	G	T	C	A	A	630
AS	K	I	R	A	N	E	L	D	V	L	K	K	L	V	F	222	
gp190n	A	T	C	T	A	T	A	A		C	T	A	A	C	T	G	
gp190s	A	A	G	A	T	C	A	G	A	T	T	G	A	C	G	T	675




FIG. 3F

AS	G	Y	R	K	P	L	D	N	I	K	D	N	V	G	K	237
gp190n			A	A	A	AT	A		T	T	A	T		A		
gp190s			GG	AT	AT	CG	CA	GC	CT	CT	CG	CA	CA	AT	CA	720
AS	M	E	D	Y	I	K	K	N	K	K	T	I	E	N	I	252
gp190n					C			A		A	A	A	A	T	A	
gp190s			AT	CG	AG	AT	T	A	T	A	AA	GA	GA	AT	AA	765
AS	N	E	L	I	E	E	S	K	K	T	I	D	K	N	K	267
gp190n			T	AT	A	T		AG	T	G	A	A	T	T		
gp190s			A	AC	GA	CT	GA	GA	AT	CC	AA	AA	GA	CA	CA	810
AS	N	A	T	K	E	E	E	K	K	K	L	Y	Q	A	Q	282
gp190n				T	A	A			A	A	A	A		A	T	
gp190s			A	AT	GA	CA	CA	GA	GA	GA	AA	AA	GA	AG	TT	855
AS	Y	D	L	S	I	Y	N	K	Q	L	E	E	A	H	N	297
gp190n			T	T	T	T	C	T		AT	A		A		T	
gp190s			T	AC	CA	CT	TA	TA	CA	CA	AA	CA	AG	CT	TA	900




FIG. 3G

AS	L I S V L E K R I D T L K K N	312
gp190n	T A A TT A A A T T TT A A A	
gp190s	CTCATCAGCGTACTGGAGAAGCGCATAGACACCCCTCAAGAAGAAT	945
AS	E N I K E L L D K I N E I K N	327
gp190n	C T G T A T T A A	
gp190s	GAAATATCAAGAAGTCTCGACAAGATTAAATGAAATTAAAGAAT	990
AS	P P P A N S G N T P N T L L D	342
gp190n	C A G T A T A A T T C T T	
gp190s	CCTCCGCCAGCCAACCTCTGGGAACACCCCTAACACGCTGCTGGAC	1035
AS	K N K K I E E H E K E I K E I	357
gp190n	A A C A A A A A A A T	
gp190s	AAGAACAAGAAGATAGAGGAGCAGAGAAAGAGATCAAAGAGATC	1080
AS	A K T I K F N I D S L F T D P	372
gp190n	T A T T T AG T A A	
gp190s	GCCAAAACCATTAAGTTCAACATAGATTCTCTCTTTACTGATCCC	1125




FIG. 3H

AS	L E L E Y Y L R E K N I D	387
gp190n	AT A A T A A A A T T	
gp190s	CTTGAGCTGGAGTACTTGGAGAGAGAATAAGAAATATAGAC	1170
AS	I S A K V E T K E S T E P N E	402
gp190n	AAGT A G T A T C	
gp190s	ATCTCCGCCAAAGTCGAGACAAAGGAATCAACCGAACCTAATGAA	1215
AS	Y P N G V T Y P L S Y N D I N	417
gp190n	A A T T T A T	
gp190s	TATCCCAATGGTGTGACGTACCCCTCTGTCTTATAACGATATCAAC	1260
AS	N A L N E L N S F G D L I N P	432
gp190n	T T A T A T TCT T T A T A	
gp190s	AACGCTCTCAACGAGCTCAATAGCTTCCGGTGACTTTCGATTAAACCCC	1305
AS	F D Y T K E P S K N I Y T D N	447
gp190n	T A AAG A C A T T T	
gp190s	TTCGATTATACGAAAGAACCCCTCTAAGAAATATCTACACAGACAAT	1350




FIG. 3I

AS	E R K K F I N E I K E K I K I	462
gp190n	A A A C A T T A A T A	
gp190s	GAGAGAAAGAGTTTATCAACGAAATCAAGGAGAGATCAAAATT	1395
AS	E K K K I E S D K K S Y E D R	477
gp190n	A A A ATC T A TC A A	
gp190s	GAGAAGAGAAATTTGAGAGTGACAAAGAAAGTTACGAAGACCGC	1440
AS	S K S L N D I T K E Y E K L L	492
gp190n	TCT GTC T T A A A AT A T	
gp190s	AGCAAAAGTCTAAACGATATCACTAAAGAGTATGAAAAGCTGCTG	1485
AS	N E I Y D S K F N N N I D L T	507
gp190n	T A T AG T T A TT A T	
gp190s	AACGAGATCTATGATTCCAAATTCAACAATAACATCGACCTGACC	1530
AS	N F E K M M G K R Y S Y K V E	522
gp190n	T A T A A T A T T	
gp190s	AACTTCGAGAAATGATGGGAAACGGTACTCTTACAAAGTGGAG	1575





FIG. 3J

AS	K	L	T	H	H	N	T	F	A	S	Y	E	N	S	K	537
gp190n			T													
gp190s	AAACTGACACACCATATAACCTTTGCATCCTATGAGAATTCTAAG															1620
AS	H	N	L	E	K	L	T	K	A	L	K	Y	M	E	D	552
gp190n					A	T	A	A							A	
gp190s	CATAATCTTGAGAAGCTCACCAAGCTCTTAAGTATATGAGGAC															1665
AS	Y	S	L	R	N	I	V	V	E	K	E	L	K	Y	Y	567
gp190n			T	A	T	A	A	T	A							
gp190s	TATTCTCTGCGGAACATTGTTGTGGAGAAAGAACTAAAGTATTAC															1710
AS	K	N	L	I	S	K	I	E	N	E	I	E	T	L	V	582
gp190n			A	T	A	C	A	A		T		T	A	A	T	
gp190s	AAGAATCTCATAAGTAAGATCGAAACGAGATCGAGACGCTTGTT															1755
AS	E	N	I	K	K	D	E	E	Q	L	F	E	K	K	I	597
gp190n			A	T	A	A				C	T		A	A	A	
gp190s	GAGAACATTAAAGAAGGATGAAGAACAGTTGTTTGAGAAGAAGATT															1800





AS	T K D E N K P D E K I L E V S	612
gp190n	T A A T T A A A T	
gp190s	ACAAAGACGAAAATAAACCATGATGAGAAAGATCCTGGAGGTCTCC	1845
AS	D I V K V Q V Q K V L L M N K	627
gp190n	C A A T A A T T A T A	
gp190s	GATATTGTTAAAGTCCAAGTGCAGAAAGGTGCTCCTCATGAACAAG	1890
AS	I D E L K K T Q L I L K N V E	642
gp190n	C T A A T G T A A T A A	
gp190s	ATTGATGAACTCAAGAAAGACTCAACTCATTTCTGAAGAACGTGGAG	1935
AS	L K H N I H V P N S Y K Q E N	657
gp190n	T C T C C A A A	
gp190s	TTAAACATAATATACATGTGCCGAATAGTTATAAGCAGAGAAT	1980
AS	K Q E P Y Y L I V L K K E I D	672
gp190n	A T T T A T G T G A A T T	
gp190s	AAGCAGGAACCATACTACCTCATCTGTACTCAAGAAAGAGATAGAC	2025




FIG. 3L

AS	K L K V F M P K V E S L I N E	687
gp190n	T A T G A ATCAT A T	
gp190s	AAACTGAAAGTGTTTCATGCCCCAAAGTCGAGAGCCTGATCAACGAA	2070
AS	E K K N I K T E G Q S D N S E	702
gp190n	A A A A A T A G T G A	
gp190s	GAGAAGAAGAACATTAAAACTGAAGGACAGTCAGATAACTCCGAG	2115
AS	P S T E G E I T G Q A T T K P	717
gp190n	A A C A A A A T A A T	
gp190s	CCTTCCACAGAAGGAGAGATAACCGGACAGGCTACCAAGCCC	2160
AS	G Q Q A G S A L E G D S V Q A	732
gp190n	A A A A T A A TCA A A	
gp190s	GGACAACAGGCCGGTTCAGCTCTCGAAGGCGATAGCGTCAAGCT	2205
AS	Q A Q E Q K Q A Q P P V P	747
gp190n	A A A A A A A A A	
gp190s	CAAGCACAGAGCAGAAGCAGGCACAGCCTCCAGTGCCAGTGCCC	2250




FIG. 3M

AS	V	P	E	A	K	A	Q	V	P	T	P	P	A	P	V	762	
gp190n	A	A	A	A	A	C	A						A	A	A		
gp190s	G	T	C	C	A	G	G	C	T	A	A	G	T	C	C	T	2295
AS	N	N	K	T	E	N	V	S	K	L	D	Y	L	E	K	777	
gp190n	T	A	T	A		T	T	C		T	A	T	T	A	A		
gp190s	A	A	T	A	C	C	G	A	A	T	G	T	C	A	G	A	2340
AS	L	Y	E	F	L	N	T	S	Y	I	C	H	K	Y	I	792	
gp190n	T	A	A	T	T	A		T	A	T	A	T			T		
gp190s	C	T	C	T	A	T	G	A	T	A	C	A	T	C	C	T	2385
AS	L	V	S	H	S	T	M	N	E	K	I	L	K	Q	Y	807	
gp190n	T	G	T	A	T	C	A		A		A	T	A	A	T		
gp190s	C	T	C	G	T	C	T	C	A	G	C	A	T	A	T	A	2430
AS	K	I	T	K	E	E	S	K	L	S	S	C	D	P		822	
gp190n	A	T	A		G	A	A	C	T	A	A	G	T	A			
gp190s	A	A	G	A	T	A	C	C	A	G	A	G	T	A	A	A	2475






FIG. 3N


AS	L D L L F N I Q N N I P V M Y	837
gp190n	T A T A T A A T A T A	
gp190S	CTGGACCTGCTGTTCAATATCCAGAACAACATTCCTGTTATGTAT	2520
AS	S M F D S L N N S L S Q L F M	852
gp190n	T T A A G T A A A T	
gp190S	TCTATGTTTCGATAGCCTCAACAATTCCTCTCTCAACTGTTTCATG	2565
AS	E I Y E K E M V C N L Y K L K	867
gp190n	A T A A A T T T A T G	
gp190S	GAGATATATGAGAAGGAGATGGTCTGCAACCTGTATAAACTCAAA	2610
AS	D N D K I K N L L E E A K K V	882
gp190n	T T A A T T A T A G A A A	
gp190S	GACAACGACAAGATTAAAGAACCTTCTGAGGAAGCTAAGAAGGTC	2655
AS	S T S V K T L S S S S M Q P L	897
gp190n	A A T A A G T T C A T A	
gp190S	TCCACCTCTGTTAAAACTCTCTCTTCCAGCTCCATGCAACCACTG	2700




AS	S L T P Q D K P E V S A N D D	912
gp190n	AT A G T A A T A T T T	
gp190S	TCTCTCACACCTCAAGACAAGCCCGAAGTGAGCGCTAACGACGAC	2745
AS	T S H S T N L N N S L K L F E	927
gp190n	A A T T A TT G TAGTT A T A A	
gp190S	ACCTCTCACTCGACC AACCTTAAATAACTCACTGA AACTGT TTGAG	2790
AS	N I L S L G K N K N I Y Q E L	942
gp190n	AT AG T A A C A T A T A T A	
gp190S	AACATCCTGTCTCTCGGCAAGAATAAGAACATCTACCAAGAACTT	2835
AS	I G Q K S S E N F Y E K I L K	957
gp190n	A T A AGTAGT A T T A T A	
gp190S	ATTGGACAGAAATCGTCCGAGAACTTCTACGAGAAGATACTGAAA	2880
AS	D S D T F Y N E S F T N F V K	972
gp190n	T T T T ATCT T A T T A	
gp190S	GACAGCGACACATTCTATAACGAGAGCTTCACCTAACTTCGTGAAA	2925



AS gp190n	S K A D D I N S L N D E S K R	987
gp190s	TCTAAAGCCGATGATATCAACTCTCTTAACGATGAATCTAAACGT	2970
AS gp190n	K K L E E D I N K L K K T L Q	1002
gp190s	AAGAAGCTGGAAGAGGACATCAATAAGCTGAAGAAGACACTGCAA	3015
AS gp190n	L S F D L Y N K Y K L K L E R	1017
gp190s	CTGAGCTTCGACCTGTACAACAAGTACAAACTGAAACTGGAGAGA	3060
AS gp190n	L F D K K K T V G K Y K M Q I	1032
gp190s	CTCTTCGACAAGAAGACAGACAGTCGGCAAGTATAAGATGCAGATC	3105
AS gp190n	K K L T L L K E Q L E S K L N	1047
gp190s	AAGAAGTTGACTCTGCTCAAGGAGCAGCTTGAAGCAAACTCAAC	3150




AS	S L N N P K H V L Q N F S V F	1062
gp190n	T T C A G T T A A T T T T	
gp190s	TCACTGAACAATCCGAAACACGTACTGCAGAACTTCTCAGTGTTCTC	3195
AS	F N K K E A E I A E T E N T	1077
gp190n	T A A A T A A A T A A	
gp190s	TTCAACAAGAAGGAAGCCGAGATCGCCGAGACAGAGAACAACACT	3240
AS	L E N T K I L L K H Y K G L V	1092
gp190n	T A A A A AT AT G T T A T T	
gp190s	CTGGAGAACACCAAGATTCTTCTCAAAACACTACAAAGGCCCTCGTC	3285
AS	K Y Y N G E S S P L K T L S E	1107
gp190n	A T A A AT A A T AAGT A	
gp190s	AAGTATTATAATGGCGAGTCTTCTCCTCTGAAGACTCTCTCCGAG	3330
AS	E S I Q T E D N Y A S L E N F	1122
gp190n	ATCA T A A A T T T A A T	
gp190s	GAGAGCATCCAGACCGAGGATAACTACGCCAGCCTCGAGAACTTC	3375



AS	K V L S K L E G K L K D N L N	1137
gp190n	A AT AAG AT A A AT A T TT A T	
gp190s	AAGTCCCTGTCTAAGCTCGAAGGCAAGCTGAAGGACAACCTGAAC	3420
AS	L E K K K L S Y L S S G L H H	1152
gp190n	T A A A AT ATCA T A A T A T	
gp190s	CTGGAGAAGAAGCTCAGCTACCTCTCTAGCGGACTGCATCAC	3465
AS	L I A E L K E V I K N K N Y T	1167
gp190n	T A T T AT A A A A A T A T T A	
gp190s	CTGATCGCCGAGCTCAAGGAAAGTCATTAAAGAACAAAGAACTACACC	3510
AS	G N S P S E N N T D V N N A L	1182
gp190n	T TCT T A G T T C T T A	
gp190s	GGCAATAGCCCCAAGCGAGAATAATACAGACGTGAATAACGCACGTG	3555
AS	E S Y K K F L P E G T D V A T	1197
gp190n	A A T C A T A A	
gp190s	GAATCTACAAGAAGTTCTTGCCCTGAAGGAACAGATGTCGCCACT	3600





AS	V V S E S G S D T L E Q S Q P	1212
gp190n	T AAG AG A T A A AAG A	
gp190S	GTGGTGTCTGAATCTGGCTCCGACACACTGGAGCAGTCTCAACCT	3645
AS	K K P A S T H V G A E S N T I	1227
gp190n	A A A A A A T C A	
gp190S	AAGAAGCCTGCATCTACTCATGTCCGAGCCGAGTCACAATAACAATT	3690
AS	T T S Q N V D D E V D D V I I	1242
gp190n	A A A T T A A A A	
gp190S	ACCACATCTCAGAACGTCGACGATGAGGTCCGATGACGTCAATCATTT	3735
AS	V P I F G E S E E D Y D D L G	1257
gp190n	A A T A ATC A A T T TT A A	
gp190S	GTGCCATATCTTCGGCGAGAGCGAGGAGGACTACGATGACCTCGGC	3780
AS	Q V V T G E A V T P S V I D N	1272
gp190n	A A A A A A A	
gp190S	CAGGTGTCACCGGTGAGGCTGTCACTCCTTCCGTGATGATAAC	3825




FIG. 3T

AS	I	L	S	K	I	E	N	E	Y	E	V	L	Y	L	K	1287
gp190n	A	T	T	T	A	T	T	T	G	T	T	A	T	A		
gp190s	ATTCTGTCCAAAATCGAGAACGAATACGGAAGTGCTCTATCTGAAA	3870														
AS	P	L	A	G	V	Y	R	S	L	K	K	Q	L	E	N	1302
gp190n	T	A	T	T	AAG	T	A	A	AT	A	A					
gp190s	CCTCTGGCAGGCGTCTATAGTCTCTCAAGAAACAGCTGGAGAAT	3915														
AS	N	V	M	T	F	N	V	N	V	K	D	I	L	N	S	1317
gp190n	T	A	T	T	T	T	T	T	T	T	T	A	TTCA			
gp190s	AACGTGATGACCTTCAATGTCAACGTGAAGGACATTTCTGAACAGC	3960														
AS	R	F	N	K	R	E	N	F	K	N	V	L	E	S	D	1332
gp190n	A	AC	T		A	T	T	A	ATCA	T						
gp190s	CGCTTTAATAAGAGAGAGAAAATTTCAGAAACGCTCTTGGAGAGCGAC	4005														
AS	L	I	P	Y	K	D	L	T	S	S	N	Y	V	V	K	1347
gp190n	A	A	TT	A	A	AAG	T	T								
gp190s	TTGATTCCCTATAAAGACCTGACCTCCTCTAACTACGTTGTCAAG	4050														




FIG. 3U

AS	D	P	Y	K	F	L	N	K	E	K	R	D	K	F	L	1362
gp190n	T	A	T	T	A	A	A	A	A	A	A	A	A	A	CT A	
gp190S	G	A	C	C	A	T	A	C	A	A	G	T	T	C	C	4095
	S	S	Y	N	Y	I	K	D	S	I	D	T	D	I	N	
AS	S	S	Y	N	Y	I	K	D	S	I	D	T	D	I	N	1377
gp190n	A	G	C	T	T	T	T	A	A	A	T	G	A	A	A	
gp190S	T	C	T	A	G	T	T	A	C	A	A	G	A	C	T	4140
	F	A	N	D	V	L	G	Y	Y	K	I	L	S	E	K	
AS	F	A	N	D	V	L	G	Y	Y	K	I	L	S	E	K	1392
gp190n	T	A	T	T	A	T	A	T	A	T	A	T	A	T	A	
gp190S	T	T	C	G	C	T	A	A	T	T	A	T	A	T	A	4185
	Y	K	S	D	L	D	S	I	K	K	Y	I	N	D	K	
AS	Y	K	S	D	L	D	S	I	K	K	Y	I	N	D	K	1407
gp190n	T	A	A	T	T	A	T	A	A	A	A	A	A	A	C A	
gp190S	T	A	C	A	G	T	C	T	G	A	C	T	T	A	A	4230
	Q	G	E	N	E	K	Y	L	P	F	L	N	N	I	E	
AS	Q	G	E	N	E	K	Y	L	P	F	L	N	N	I	E	1422
gp190n	T	A	G	C	T	T	T	A	C	T	T	A	C	T	T G	
gp190S	C	A	A	G	G	A	A	T	A	T	C	T	G	A	A	4275




FIG. 3V

AS	T	L	Y	K	T	V	N	D	K	I	D	L	F	V	I	1437
gp190n	T	A	T	A	T	T	T	T	T	T	T	T	T	A	T	
gp190s	ACCCTGTACAAGACAGTGAACGACGACAAATCGACCTCTTCGTAATT	4320														
AS	H	L	E	A	K	V	L	N	Y	T	Y	E	K	S	N	1452
gp190n	TTA A A A T A T A T A T ATCA C															
gp190s	CACCTGGAGGCCAAGTGCTCAACTATACTTACGAGAGAGCAAT	4365														
AS	V	E	V	K	I	K	E	L	N	Y	L	K	T	I	Q	1467
gp190n	A A A A T T T A T															
gp190s	GTGGAAGTTAAATCAAGGAGCTGAACCTACCTCAAAACAATCCAA	4410														
AS	D	K	L	A	D	F	K	K	N	N	N	F	V	G	I	1482
gp190n	AT T A															
gp190s	GACAAGCTGGCAGATTTCAGAAGAAAATAACAATTTCGTCGGAATT	4455														
AS	A	D	L	S	T	D	Y	N	H	N	N	L	L	T	K	1497
gp190n	T TT A A A T T CT AT A															
gp190s	GCAGACCTGTCTACCGATTATAACCACACAACAATCTCCTGACCAAG	4500														




FIG. 3W

AS	F L S T G M V F E N L A K T V	1512
gp190n	C TAGT A T T T T C T	
gp190s	TTTCTGTCCACTGGCATGGTGTTCGAAACCTCGCCAAACAGTG	4545
AS	L S N L L D G N L Q G M L N I	1527
gp190n	T ATCT T A T A T A T A T	
gp190s	CTGAGCAATCTGCTCGACGGCAACCTGCAGGGCATGCTGAACATC	4590
AS	S Q H Q C V K K Q C P Q N S G	1542
gp190n	A A A A A A T A A TCT A	
gp190s	TCCCAGCACCAATGCGTGAAGAAACAGTGCCCCCAGAAATAGCGGC	4635
AS	C F R H L D E R E E C K C L L	1557
gp190n	A T A T AA A A T A T AT A	
gp190s	TGTTTCAGGCATCTGGACGAGCGCGAAGAGTGCAAGTGTCTCCTG	4680
AS	N Y K Q E G D K C V E N P N P	1572
gp190n	T T A T A T T	
gp190s	AACTACAAACAAGAGGAGATAAGTGCCTGGAGAACCCAAACCTT	4725




FIG. 3X

AS	T C N E N N G G C D A D A K C	1587
gp190n	T T C T A T A C T	
gp190S	ACCTGCAATGAAACAATGGCGGTGTGACGCCGATGCTAAATGC	4770
AS	T E E D S G S N G K K I T C E	1602
gp190n	A TTCA TAGC T A	
gp190S	ACCGAGGAAGACAGCGGCTCTAACGGAAAGAAATCACATGCCGAG	4815
AS	C T K P D S Y P L F D G I F C	1617
gp190n	A T T T T T T C	
gp190S	TGTACTAAGCCCGACTCCTATCCACTCTTCGACGGGATTTTTTGC	4860
AS	S S S N F L G I F F L L I L M	1632
gp190n	AGTTC C T A A A C A T A T A A	
gp190S	TCCAGCTCTAATTTCCCTGGGCATCTTCTTCTCCTGCTGATCCTCATG	4905
AS	L I L Y S F I * *	1639
gp190n	T A A T A T T	
gp190S	CTGATCCTGTACAGCTTCATCTAATAGATCGATGG	4940
	stop codon Cla I	

FIG. 3Y

	N'-terminus	C'-terminus
<b>gp190s1 Sequence</b>		
DNA Sequence	GC <u>ACGCGTATGAAATC</u> ----- AGCTCTAATTAAATAGGCGGCCGCATCGATGGC	
AA Sequence	Mlu I Met Lys Ile ----- Ser Ser Asn stop codon Not I Cla I	
AA Position	1 2 3 ----- 1619 1620 1621	
<b>gp190s2 Sequence</b>		
DNA Sequence	GC <u>GGATCCGTGACCCAC</u> ----- AGCTCTAATTAAATAGGCGGCCGCATCGATGGC	
AA Sequence	BamH I Val Thr His ----- Ser Ser Asn stop codon Not I Cla I	
AA Position	20 21 22 ----- 1619 1620 1621	







FIG. 4A









FIG. 4B







FIG. 5A







FIG. 5B







FIG. 6A






FIG. 6B

